antibodies .- online.com





TRIM13 Protein (AA 1-407) (Strep Tag)



Image



Overview

Quantity:	1 mg
Target:	TRIM13
Protein Characteristics:	AA 1-407
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRIM13 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MELLEEDLTC PICCSLFDDP RVLPCSHNFC KKCLEGILEG SVRNSLWRPA PFKCPTCRKE TSATGINSLQ VNYSLKGIVE KYNKIKISPK MPVCKGHLGQ PLNIFCLTDM QLICGICATR GEHTKHVFCS IEDAYAQERD AFESLFQSFE TWRRGDALSR LDTLETSKRK SLQLLTKDSD KVKEFFEKLQ HTLDQKKNEI LSDFETMKLA VMQAYDPEIN KLNTILQEQR MAFNIAEAFK DVSEPIVFLQ QMQEFREKIK VIKETPLPPS NLPASPLMKN FDTSQWEDIK LVDVDKLSLP ODTGTFISKI PWSFYKLFLL ILLLGLVIVF GPTMFLEWSL FDDLATWKGC LSNFSSYLTK TADFIEQSVF YWEQVTDGFF IFNERFKNFT LVVLNNVAEF VCKYKLL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Product Details

Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

Target Details	
Target:	TRIM13
Alternative Name:	TRIM13 (TRIM13 Products)
Background:	E3 ubiquitin-protein ligase TRIM13 (EC 2.3.2.27) (B-cell chronic lymphocytic leukemia tumor
	suppressor Leu5) (Leukemia-associated protein 5) (Putative tumor suppressor RFP2) (RING
	finger protein 77) (RING-type E3 ubiquitin transferase TRIM13) (Ret finger protein 2) (Tripartite
	motif-containing protein 13),FUNCTION: Endoplasmic reticulum (ER) membrane anchored E3
	ligase involved in the retrotranslocation and turnover of membrane and secretory proteins from
	the ER through a set of processes named ER-associated degradation (ERAD). This process act
	on misfolded proteins as well as in the regulated degradation of correctly folded proteins.
	Enhances ionizing radiation-induced p53/TP53 stability and apoptosis via ubiquitinating MDM2
	and AKT1 and decreasing AKT1 kinase activity through MDM2 and AKT1 proteasomal
	degradation. Regulates ER stress-induced autophagy, and may act as a tumor suppressor
	(PubMed:22178386). Also plays a role in innate immune response by stimulating NF-kappa-B
	activity in the TLR2 signaling pathway. Ubiquitinates TRAF6 via the 'Lys-29'-linked
	polyubiquitination chain resulting in NF-kappa-B activation (PubMed:28087809). Participates a
	well in T-cell receptor-mediated NF-kappa-B activation (PubMed:25088585). In the presence of
	TNF, modulates the IKK complex by regulating IKBKG/NEMO ubiquitination leading to the
	repression of NF-kappa-B (PubMed:25152375). {ECO:0000269 PubMed:17314412,
	ECO:0000269 PubMed:21333377, ECO:0000269 PubMed:22178386,
	ECO:0000269 PubMed:25088585, ECO:0000269 PubMed:25152375,
	ECO:0000269 PubMed:28087809}.
Molecular Weight:	47.0 kDa
UniProt:	060858
UniProt:	U6U858

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process