antibodies .- online.com





Transferrin Receptor Protein (AA 101-760) (rho-1D4 tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	Transferrin Receptor (TFRC)
Protein Characteristics:	AA 101-760
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Transferrin Receptor protein is labelled with rho-1D4 tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

LAGTESPVRE EPGEDFPAAR RLYWDDLKRK LSEKLDSTDF TGTIKLLNEN SYVPREAGSQ
KDENLALYVE NQFREFKLSK VWRDQHFVKI QVKDSAQNSV IIVDKNGRLV YLVENPGGYV
AYSKAATVTG KLVHANFGTK KDFEDLYTPV NGSIVIVRAG KITFAEKVAN AESLNAIGVL
IYMDQTKFPI VNAELSFFGH AHLGTGDPYT PGFPSFNHTQ FPPSRSSGLP NIPVQTISRA
AAEKLFGNME GDCPSDWKTD STCRMVTSES KNVKLTVSNV LKEIKILNIF GVIKGFVEPD
HYVVVGAQRD AWGPGAAKSG VGTALLLKLA QMFSDMVLKD GFQPSRSIIF ASWSAGDFGS
VGATEWLEGY LSSLHLKAFT YINLDKAVLG TSNFKVSASP LLYTLIEKTM QNVKHPVTGQ
FLYQDSNWAS KVEKLTLDNA AFPFLAYSGI PAVSFCFCED TDYPYLGTTM DTYKELIERI
PELNKVARAA AEVAGQFVIK LTHDVELNLD YERYNSQLLS FVRDLNQYRA DIKEMGLSLQ
WLYSARGDFF RATSRLTTDF GNAEKTDRFV MKKLNDRVMR VEYHFLSPYV SPKESPFRHV
FWGSGSHTLP ALLENLKLRK QNNGAFNETL FRNQLALATW TIQGAANALS GDVWDIDNEF

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a

special request, please contact us.

Characteristics:

- · Made in Germany from design to production by highly experienced protein experts.
- Human TFRC Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin-free.

Product Details	
Grade:	Crystallography grade
Target Details	
Target:	Transferrin Receptor (TFRC)
Alternative Name:	TFRC (TFRC Products)
Background:	Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. {ECO:0000250, ECO:0000269 PubMed:3568132}., (Microbial infection) Acts as a receptor for new-world arenaviruses: Guanarito, Junin and Machupo virus. {ECO:0000269 PubMed:17287727, ECO:0000269 PubMed:18268337}.
Molecular Weight:	75.0 kDa Including tag.
UniProt:	P02786
Pathways:	Transition Metal Ion Homeostasis
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)