

## Datasheet for ABIN3112634

# IL-8 Protein (AA 27-99) (His tag)



### Overview

Quantity:	2 mg
Target:	IL-8 (IL8)
Protein Characteristics:	AA 27-99
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL-8 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)
Product Details	
Sequence:	RSAKELRCQC IKTYSKPFHP KFIKELRVIE SGPHCANTEI IVKLSDGREL CLDPKENWVQ
	RVVEKFLKRA ENS
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	Made in Germany - from design to production - by highly experienced protein experts.
	Human CXCL8 Protein (raised in E. Coli) purified by multi-step, protein-specific process to
	<ul><li>ensure crystallization grade.</li><li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul>
	otate of the art algorithm asea for plasmia design (some synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein

cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

## **Target Details**

Target:	IL-8 (IL8)
Alternative Name:	CXCL8 (IL8 Products)
Background:	IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes.
	It is also involved in neutrophil activation. It is released from several cell types in response to an inflammatory stimulus. IL-8(6-77) has a 5-10-fold higher activity on neutrophil activation, IL-8(5-
	77) has increased activity on neutrophil activation and IL-8(7-77) has a higher affinity to
	receptors CXCR1 and CXCR2 as compared to IL-8(1-77), respectively.

## **Target Details**

Expiry Date:

l arget Details	
	{ECO:0000269 PubMed:11978786, ECO:0000269 PubMed:2145175,
	ECO:0000269 PubMed:2212672}.
Molecular Weight:	9.5 kDa Including tag.
UniProt:	P10145
Pathways:	TLR Signaling, Cellular Response to Molecule of Bacterial Origin, Regulation of G-Protein
	Coupled Receptor Protein Signaling, ER-Nucleus Signaling, Hepatitis C, Autophagy
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
E in . D . t	

Unlimited (if stored properly)