antibodies -online.com





CD79a Protein (AA 33-226) (rho-1D4 tag)



Overview

Overview	
Quantity:	1 mg
Target:	CD79a (CD79A)
Protein Characteristics:	AA 33-226
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD79a protein is labelled with rho-1D4 tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)
Product Details	
Sequence:	LWMHKVPASL MVSLGEDAHF QCPHNSSNNA NVTWWRVLHG NYTWPPEFLG PGEDPNGTLI
	IQNVNKSHGG IYVCRVQEGN ESYQQSCGTY LRVRQPPPRP FLDMGEGTKN RIITAEGIIL
	LFCAVVPGTL LLFRKRWQNE KLGLDAGDEY EDENLYEGLN LDDCSMYEDI SRGLQGTYQD
	VGSLNIGDVQ LEKP
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	Made in Germany - from design to production - by highly experienced protein experts.
	 Human CD79A Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our

experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin-free.

Grade: Crystallography grade

Target Details

Target:	CD79a (CD79A)
Alternative Name:	CD79A (CD79A Products)
Background:	Required in cooperation with CD79B for initiation of the signal transduction cascade activated

by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization		
of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR		
surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK		
autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and		
allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-		
family tyrosine kinases. Represses BCR signaling during development of immature B-cells.		
{ECO:0000269 PubMed:8617796, ECO:0000269 PubMed:9057631}.		

In addition to the applications listed above we expect the protein to work for functional studies

Molecular Weight:	23.0 kDa Including tag.
UniProt:	P11912
Pathways:	BCR Signaling

Application Details

Application Notes:

1.1	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)