antibodies -online.com





Datasheet for ABIN3113031

XBP1 Protein (AA 1-261) (Strep Tag)



Overview

Quantity:	1 mg
Target:	XBP1
Protein Characteristics:	AA 1-261
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This XBP1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence:

MVVVAAAPNP ADGTPKVLLL SGQPASAAGA PAGQALPLMV PAQRGASPEA ASGGLPQARK RQRLTHLSPE EKALRRKLKN RVAAQTARDR KKARMSELEQ QVVDLEEENQ KLLLENQLLR EKTHGLVVEN QELRQRLGMD ALVAEEEAEA KGNEVRPVAG SAESAALRLR APLQQVQAQL SPLQNISPWI LAVLTLQIQS LISCWAFWTT WTQSCSSNAL PQSLPAWRSS QRSTQKDPVP YQPPFLCQWG RHQPSWKPLM N

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Target Details

- Target Betane	
Target:	XBP1
Alternative Name:	XBP1 (XBP1 Products)
Background:	X-box-binding protein 1 (XBP-1) (Tax-responsive element-binding protein 5) (TREB-5) [Cleaved
	into: X-box-binding protein 1, cytoplasmic form, X-box-binding protein 1, luminal
	form],FUNCTION: Functions as a transcription factor during endoplasmic reticulum (ER) stress
	by regulating the unfolded protein response (UPR). Required for cardiac myogenesis and
	hepatogenesis during embryonic development, and the development of secretory tissues such
	as exocrine pancreas and salivary gland (By similarity). Involved in terminal differentiation of B
	lymphocytes to plasma cells and production of immunoglobulins (PubMed:11460154).
	Modulates the cellular response to ER stress in a PIK3R-dependent manner
	(PubMed:20348923). Binds to the cis-acting X box present in the promoter regions of major
	histocompatibility complex class II genes (PubMed:8349596). Involved in VEGF-induced
	endothelial cell (EC) proliferation and retinal blood vessel formation during embryonic
	development but also for angiogenesis in adult tissues under ischemic conditions. Functions
	also as a major regulator of the UPR in obesity-induced insulin resistance and type 2 diabetes
	for the management of obesity and diabetes prevention (By similarity).
	{ECO:0000250 UniProtKB:035426, ECO:0000269 PubMed:11460154,
	ECO:0000269 PubMed:20348923, ECO:0000269 PubMed:8349596}., FUNCTION: [Isoform 1]:
	Plays a role in the unconventional cytoplasmic splicing processing of its own mRNA triggered
	by the endoplasmic reticulum (ER) transmembrane endoribonuclease ERN1: upon ER stress,
	the emerging XBP1 polypeptide chain, as part of a mRNA-ribosome-nascent chain (R-RNC)
	complex, cotranslationally recruits its own unprocessed mRNA through transient docking to the
	ER membrane and translational pausing, therefore facilitating efficient IRE1-mediated XBP1
	mRNA isoform 2 production (PubMed:19394296, PubMed:21233347). In endothelial cells (EC),
	associated with KDR, promotes IRE1-mediated XBP1 mRNA isoform 2 productions in a
	vascular endothelial growth factor (VEGF)-dependent manner, leading to EC proliferation and
	angiogenesis (PubMed:23529610). Functions as a negative feed-back regulator of the potent
	transcription factor XBP1 isoform 2 protein levels through proteasome-mediated degradation,
	thus preventing the constitutive activation of the ER stress response signaling pathway
	(PubMed:16461360, PubMed:25239945). Inhibits the transactivation activity of XBP1 isoform 2
	in myeloma cells (By similarity). Acts as a weak transcriptional factor (PubMed:8657566).
	Together with HDAC3, contributes to the activation of NFE2L2-mediated HMOX1 transcription
	factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to EC
	survival under disturbed flow/oxidative stress (PubMed:25190803). Binds to the ER stress

response element (ERSE) upon ER stress (PubMed:11779464). Binds to the consensus 5'-

GATGACGTG[TG]N(3)[AT]T-3' sequence related to cAMP responsive element (CRE)-like sequences (PubMed:8657566). Binds the Tax-responsive element (TRE) present in the long terminal repeat (LTR) of T-cell leukemia virus type 1 (HTLV-I) and to the TPA response elements (TRE) (PubMed:2321018, PubMed:2196176, PubMed:1903538, PubMed:8657566). Associates preferentially to the HDAC3 gene promoter region in a static flow-dependent manner (PubMed:25190803). Binds to the CDH5/VE-cadherin gene promoter region (PubMed:19416856). (ECO:0000250|UniProtKB:035426, ECO:0000269|PubMed:11779464, ECO:0000269|PubMed:16461360, ECO:0000269|PubMed:1903538, ECO:0000269|PubMed:19394296, ECO:0000269|PubMed:19416856, ECO:0000269|PubMed:21233347, ECO:0000269|PubMed:2196176, ECO:0000269|PubMed:2321018, ECO:0000269|PubMed:23529610, ECO:0000269|PubMed:25190803, ECO:0000269|PubMed:252339945,

ECO:0000269|PubMed:8657566}, FUNCTION: [Isoform 2]: Functions as a stress-inducible potent transcriptional activator during endoplasmic reticulum (ER) stress by inducing unfolded protein response (UPR) target genes via binding to the UPR element (UPRE). Up-regulates target genes encoding ER chaperones and ER-associated degradation (ERAD) components to enhance the capacity of productive folding and degradation mechanism, respectively, in order to maintain the homeostasis of the ER under ER stress (PubMed:11779464, PubMed:25239945). Plays a role in the production of immunoglobulins and interleukin-6 in the presence of stimuli required for plasma cell differentiation (By similarity). Induces phospholipid biosynthesis and ER expansion (PubMed:15466483). Contributes to the VEGF-induced endothelial cell (EC) growth and proliferation in a Akt/GSK-dependent and/or-independent signaling pathway, respectively, leading to beta-catenin nuclear translocation and E2F2 gene expression (PubMed:23529610). Promotes umbilical vein EC apoptosis and atherosclerotisis development in a caspase-dependent signaling pathway, and contributes to VEGF-induced EC proliferation and angiogenesis in adult tissues under ischemic conditions (PubMed:19416856, PubMed:23529610). Involved in the regulation of endostatin-induced autophagy in EC through BECN1 transcriptional activation (PubMed:23184933). Plays a role as an oncogene by promoting tumor progression: stimulates zinc finger protein SNAI1 transcription to induce epithelial-to-mesenchymal (EMT) transition, cell migration and invasion of breast cancer cells (PubMed:25280941). Involved in adipocyte differentiation by regulating lipogenic gene expression during lactation. Plays a role in the survival of both dopaminergic neurons of the substantia nigra pars compacta (SNpc), by maintaining protein homeostasis and of myeloma cells. Increases insulin sensitivity in the liver as a response to a high carbohydrate diet, resulting in improved glucose tolerance. Improves also glucose homeostasis in an ER stress- and/or insulin-independent manner through both binding and proteasome-induced degradation of the

transcription factor FOXO1, hence resulting in suppression of gluconeogenic genes expression and in a reduction of blood glucose levels. Controls the induction of de novo fatty acid synthesis in hepatocytes by regulating the expression of a subset of lipogenic genes in an ER stress- and UPR-independent manner (By similarity). Associates preferentially to the HDAC3 gene promoter region in a disturbed flow-dependent manner (PubMed:25190803). Binds to the BECN1 gene promoter region (PubMed:23184933). Binds to the CDH5/VE-cadherin gene promoter region (PubMed:19416856). Binds to the ER stress response element (ERSE) upon ER stress (PubMed:11779464). Binds to the 5'-CCACG-3' motif in the PPARG promoter (By similarity). (ECO:0000250|UniProtKB:035426, ECO:0000269|PubMed:11779464, ECO:0000269|PubMed:15466483, ECO:0000269|PubMed:19416856, ECO:0000269|PubMed:23184933, ECO:0000269|PubMed:25239945, ECO:0000269|PubMed:25190803, ECO:0000269|PubMed:25239945, ECO:0000269|PubMed:25280941}.

Molecular Weight:

28.7 kDa

UniProt:

P17861

Pathways:

ER-Nucleus Signaling, Unfolded Protein Response

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)