# antibodies .- online.com





### CCR5 Protein (AA 1-352) (Strep Tag)





Go to Product page

#### Overview

Quantity:	1 mg
Target:	CCR5
Protein Characteristics:	AA 1-352
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCR5 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

#### **Product Details**

#### Sequence:

MDYQVSSPIY DINYYTSEPC QKINVKQIAA RLLPPLYSLV FIFGFVGNML VILILINCKR
LKSMTDIYLL NLAISDLFFL LTVPFWAHYA AAQWDFGNTM CQLLTGLYFI GFFSGIFFII
LLTIDRYLAV VHAVFALKAR TVTFGVVTSV ITWVVAVFAS LPGIIFTRSQ KEGLHYTCSS
HFPYSQYQFW KNFQTLKIVI LGLVLPLLVM VICYSGILKT LLRCRNEKKR HRAVRLIFTI
MIVYFLFWAP YNIVLLLNTF QEFFGLNNCS SSNRLDQAMQ VTETLGMTHC CINPIIYAFV
GEKFRNYLLV FFQKHIAKRF CKCCSIFQQE APERASSVYT RSTGEQEISV GL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details	
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade
Target Details	
Target:	CCR5
Alternative Name:	CCR5 (CCR5 Products)
Background:	C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5) (CHEMR13) (HIV-1
	fusion coreceptor) (CD antigen CD195),FUNCTION: Receptor for a number of inflammatory CC-
	chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently
	transduces a signal by increasing the intracellular calcium ion level. May play a role in the
	control of granulocytic lineage proliferation or differentiation. Participates in T-lymphocyte
	migration to the infection site by acting as a chemotactic receptor (PubMed:30713770).
	{ECO:0000269 PubMed:10383387, ECO:0000269 PubMed:11323418,
	ECO:0000269 PubMed:30713770, ECO:0000269 PubMed:8639485,
	ECO:0000269 PubMed:8663314, ECO:0000269 PubMed:8699119}., FUNCTION: (Microbial
	infection) Acts as a coreceptor (CD4 being the primary receptor) of human immunodeficiency
	virus-1/HIV-1. {ECO:0000269 PubMed:10383387, ECO:0000269 PubMed:21763489,
	ECO:0000269 PubMed:8649511, ECO:0000269 PubMed:8649512,
	ECO:0000269 PubMed:9632396}.
Molecular Weight:	40.5 kDa
UniProt:	P51681
Pathways:	Cellular Response to Molecule of Bacterial Origin, cAMP Metabolic Process, Regulation of Cell Size
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational

modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!


Restrictions:

For Research Use only


#### Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

#### **Images**




**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process



#### **SDS-PAGE**

Image 2. C-C Chemokine Receptor Type 5 (CCR5) (AA 1-352), fraction 10 - 11



## Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 3.** C-C Chemokine Receptor Type 5 (CCR5) (AA 1-352), gel filtration Superose 6 fraction 10 - 11

Please check the product details page for more images. Overall 4 images are available for ABIN3113835.