antibodies .- online.com







IFITM3 Protein (AA 1-133) (Strep Tag)



Image



Go to Product page

()	ve	K\ /		
	\cup	1 V/	-	V۷

Quantity:	1 mg
Target:	IFITM3
Protein Characteristics:	AA 1-133
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IFITM3 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)
Product Details	

3 , 3	·
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)
Product Details	
Sequence:	MNHTVQTFFS PVNSGQPPNY EMLKEEHEVA VLGAPHNPAP PTSTVIHIRS ETSVPDHVVW
	SLFNTLFMNP CCLGFIAFAY SVKSRDRKMV GDVTGAQAYA STAKCLNIWA LILGILMTIL
	LIVIPVLIFQ AYG
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:
	Made in Germany - from design to production - by highly experienced protein experts.
	Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/5 | Product datasheet for ABIN3114264 | 04/16/2024 | Copyright antibodies-online. All rights reserved.

reported (not tested by us and not guaranteed).

• These proteins are normally active (enzymatically functional) as our customers have

correct folding and modification.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System		
	(ALiCE®):		
	1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.		
	 Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. 		
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.		
Endotoxin Level:	vel: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)		
Grade:	Crystallography grade		

Target Details

Target:	IFITM3
Alternative Name:	IFITM3 (IFITM3 Products)
Background:	Interferon-induced transmembrane protein 3 (Dispanin subfamily A member 2b) (DSPA2b)
	(Interferon-inducible protein 1-8U), FUNCTION: IFN-induced antiviral protein which disrupts
	intracellular cholesterol homeostasis. Inhibits the entry of viruses to the host cell cytoplasm by
	preventing viral fusion with cholesterol depleted endosomes. May inactivate new enveloped
	viruses which buds out of the infected cell, by letting them go out with a cholesterol depleted
	membrane. Active against multiple viruses, including influenza A virus, SARS coronaviruses
	(SARS-CoV and SARS-CoV-2), Marburg virus (MARV), Ebola virus (EBOV), Dengue virus (DNV),
	West Nile virus (WNV), human immunodeficiency virus type 1 (HIV-1), hepatitis C virus (HCV)
	and vesicular stomatitis virus (VSV) (PubMed:26354436, PubMed:33270927,
	PubMed:33239446). Can inhibit: influenza virus hemagglutinin protein-mediated viral entry,
	MARV and EBOV GP1,2-mediated viral entry, SARS-CoV and SARS-CoV-2 S protein-mediated
	viral entry and VSV G protein-mediated viral entry (PubMed:33270927). Plays a critical role in
	the structural stability and function of vacuolar ATPase (v-ATPase). Establishes physical
	contact with the v-ATPase of endosomes which is critical for proper clathrin localization and is
	also required for the function of the v-ATPase to lower the pH in phagocytic endosomes thus
	establishing an antiviral state. In hepatocytes, IFITM proteins act in a coordinated manner to
	restrict HCV infection by targeting the endocytosed HCV virion for lysosomal degradation
	(PubMed:26354436). IFITM2 and IFITM3 display anti-HCV activity that may complement the
	anti-HCV activity of IFITM1 by inhibiting the late stages of HCV entry, possibly in a coordinated
	manner by trapping the virion in the endosomal pathway and targeting it for degradation at the
	lysosome (PubMed:26354436). Exerts opposing activities on SARS-CoV-2, including
	amphipathicity-dependent restriction of virus at endosomes and amphipathicity-independent
	enhancement of infection at the plasma membrane (PubMed:33270927).
	{ECO:0000269 PubMed:20064371, ECO:0000269 PubMed:20534863,
	ECO:0000269 PubMed:20943977, ECO:0000269 PubMed:21177806,
	ECO:0000269 PubMed:21253575, ECO:0000269 PubMed:22046135,
	ECO:0000269 PubMed:22479637, ECO:0000269 PubMed:23601107,
	ECO:0000269 PubMed:26354436, ECO:0000269 PubMed:33239446,
	ECO:0000269 PubMed:33270927}.
Molecular Weight:	14.6 kDa
UniProt:	Q01628

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for	
	protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process