

Datasheet for ABIN3114328 **TAP2 Protein (AA 1-686) (Strep Tag)**



Go to Product page

| _ | | | | | | |
|---|-------|-------------|----|----|-------------|-----|
| | V | \triangle | r۱ | /1 | \triangle | Λ/ |
| | ' V ' | | ΙV | | | v v |

| Quantity: | 250 μg |
|-------------------------------|---|
| Target: | TAP2 |
| Protein Characteristics: | AA 1-686 |
| Origin: | Human |
| Source: | Cell-free protein synthesis (CFPS) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This TAP2 protein is labelled with Strep Tag. |
| Application: | Western Blotting (WB), SDS-PAGE (SDS), ELISA |

| Application: | Western Blotting (WB), SDS-PAGE (SDS), ELISA | | | |
|-----------------|---|--|--|--|
| Product Details | | | | |
| Brand: | AliCE® | | | |
| Sequence: | MRLPDLRPWT SLLLVDAALL WLLQGPLGTL LPQGLPGLWL EGTLRLGGLW GLLKLRGLLG | | | |
| | FVGTLLLPLC LATPLTVSLR ALVAGASRAP PARVASAPWS WLLVGYGAAG LSWSLWAVLS | | | |
| | PPGAQEKEQD QVNNKVLMWR LLKLSRPDLP LLVAAFFFLV LAVLGETLIP HYSGRVIDIL | | | |
| | GGDFDPHAFA SAIFFMCLFS FGSSLSAGCR GGCFTYTMSR INLRIREQLF SSLLRQDLGF | | | |
| | FQETKTGELN SRLSSDTTLM SNWLPLNANV LLRSLVKVVG LYGFMLSISP RLTLLSLLHM | | | |
| | PFTIAAEKVY NTRHQEVLRE IQDAVARAGQ VVREAVGGLQ TVRSFGAEEH EVCRYKEALE | | | |
| | QCRQLYWRRD LERALYLLVR RVLHLGVQML MLSCGLQQMQ DGELTQGSLL SFMIYQESVG | | | |
| | SYVQTLVYIY GDMLSNVGAA EKVFSYMDRQ PNLPSPGTLA PTTLQGVVKF QDVSFAYPNR | | | |
| | PDRPVLKGLT FTLRPGEVTA LVGPNGSGKS TVAALLQNLY QPTGGQVLLD EKPISQYEHC | | | |
| | YLHSQVVSVG QEPVLFSGSV RNNIAYGLQS CEDDKVMAAA QAAHADDFIQ EMEHGIYTDV | | | |
| | GEKGSQLAAG QKQRLAIARA LVRDPRVLIL DEATSALDVQ CEQALQDWNS RGDRTVLVIA | | | |

HRLQTVQRAH QILVLQEGKL QKLAQL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- · The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).

| Product Details | | | |
|-------------------|---|--|--|
| Purity: | > 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC). | | |
| Grade: | custom-made | | |
| Target Details | | | |
| Target: | TAP2 | | |
| Alternative Name: | TAP2 (TAP2 Products) | | |
| Background: | Antigen peptide transporter 2 (APT2) (EC 7.4.2.14) (ATP-binding cassette sub-family B member | | |
| | 3) (Peptide supply factor 2) (Peptide transporter PSF2) (PSF-2) (Peptide transporter TAP2) | | |
| | (Peptide transporter involved in antigen processing 2) (Really interesting new gene 11 protein) | | |
| | (RING11),FUNCTION: ABC transporter associated with antigen processing. In complex with | | |
| | TAP1 mediates unidirectional translocation of peptide antigens from cytosol to endoplasmic | | |
| | reticulum (ER) for loading onto MHC class I (MHCI) molecules (PubMed:25656091, | | |
| | PubMed:25377891). Uses the chemical energy of ATP to export peptides against the | | |
| | concentration gradient (PubMed:25377891). During the transport cycle alternates between | | |
| | 'inward-facing' state with peptide binding site facing the cytosol to 'outward-facing' state with | | |
| | peptide binding site facing the ER lumen. Peptide antigen binding to ATP-loaded TAP1-TAP2 | | |
| | induces a switch to hydrolysis-competent 'outward-facing' conformation ready for peptide | | |
| | loading onto nascent MHCI molecules. Subsequently ATP hydrolysis resets the transporter to | | |
| | the 'inward facing' state for a new cycle (PubMed:25377891, PubMed:25656091, | | |
| | PubMed:11274390). Typically transports intracellular peptide antigens of 8 to 13 amino acids | | |
| | that arise from cytosolic proteolysis via IFNG-induced immunoproteasome. Binds peptides with | | |
| | free N- and C-termini, the first three and the C-terminal residues being critical. Preferentially | | |
| | selects peptides having a highly hydrophobic residue at position 3 and hydrophobic or charged | | |
| | residues at the C-terminal anchor. Proline at position 2 has the most destabilizing effect | | |
| | (PubMed:7500034, PubMed:9256420, PubMed:11274390). As a component of the peptide | | |
| | loading complex (PLC), acts as a molecular scaffold essential for peptide-MHCI assembly and | | |
| | antigen presentation (PubMed:26611325, PubMed:1538751, PubMed:25377891). | | |
| | {ECO:0000269 PubMed:11274390, ECO:0000269 PubMed:1538751, | | |
| | ECO:0000269 PubMed:25377891, ECO:0000269 PubMed:25656091, | | |
| | ECO:0000269 PubMed:26611325, ECO:0000269 PubMed:7500034, | | |

Molecular Weight:

75.7 kDa

ECO:0000269|PubMed:9256420}.

UniProt:

Q03519

Target Details

Pathways:

Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,

Human Leukocyte Antigen (HLA) in Adaptive Immune Response

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|------------------|--|
| Buffer: | The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | 12 months |