

Datasheet for ABIN3114599
SLC13A2 Protein (AA 1-592) (Strep Tag)[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	SLC13A2
Protein Characteristics:	AA 1-592
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC13A2 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence: MATCWQALWA YRSYLIVFFV PILLPLPIL VPSKEAYCAY AILMALFWC TEALPLAVTA
LFPLILFPM GIVDASEVAV EYLKDSNLLF FGGLLVIAIV EHWNLHKRIA LRVLLIVGVR
PAPLILGFML VTAFLSMWIS NTATSAMMVP IAHAVLDQLH SSQASSNVEE GSNNPTFELQ
EPSPQKEVTK LDNGQALPVT SASSEGRAHL SQKHLHLTQC MSLCVCYSAS IGGIATLTGT
APNLVLQGGI NSLFPQNGNV VNFASWFSFA FPTMVILLLL AWLWLQILFL GFNFRKNFGI
GEKMQEQQQA AYCVIQTEHR LLGPMTFAEK AISILFVILV LLWFTREPGF FLGWGNLAFP
NAKGESMVSD GTVAIFIGII MFIIPSKFPG LTQDPENPGK LKAPLGLLDW KTVNQKMPWN
IVLLGGGYA LAKGSERSGL SEWLGKNLTP LQSVPAIA ILSLLVATF TECTSNVATT
TIFLPILASM AQACLHPLY VMLPCTLATS LAFMLPVATP PNAIVFSFGD LKVLDMARAG
FLLNIIGVLI IALAINSWG I PLFSLHSFPS WAQSNTTAQC LPSLANTTTP SP

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.

Product Details

2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

Target Details

Target:	SLC13A2
Alternative Name:	SLC13A2 (SLC13A2 Products)
Background:	<p>Solute carrier family 13 member 2 (Na⁺)/dicarboxylate cotransporter 1) (NaDC-1) (Renal sodium/dicarboxylate cotransporter),FUNCTION: Low-affinity sodium-dicarboxylate cotransporter, that mediates the entry of citric acid cycle intermediates, such as succinate, citrate, fumarate and alpha-ketoglutarate (2-oxoglutarate) into the small intestine and renal proximal tubule (PubMed:10894787, PubMed:9668069, PubMed:8967342). Transports the dicarboxylate into the cell with a probable stoichiometry of 3 Na⁺ for 1 divalent dicarboxylate, rendering the process electrogenic (PubMed:8967342, PubMed:9668069, PubMed:10894787). Citrate is transported in protonated form as a divalent anion, rather than the trivalent form which is normally found in blood (PubMed:10894787). Has a critical role in renal dicarboxylate transport (By similarity). {ECO:0000250 UniProtKB:Q9ES88, ECO:0000269 PubMed:10894787, ECO:0000269 PubMed:8967342, ECO:0000269 PubMed:9668069}.</p>
Molecular Weight:	64.4 kDa
UniProt:	Q13183

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

Application Details

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process