antibodies .- online.com





HSD17B12 Protein (AA 1-312) (rho-1D4 tag)



Image



Go to Product pag

()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	1 mg
Target:	HSD17B12
Protein Characteristics:	AA 1-312
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HSD17B12 protein is labelled with rho-1D4 tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequ	ience:
------	--------

MESALPAAGF LYWVGAGTVA YLALRISYSL FTALRVWGVG NEAGVGPGLG EWAVVTGSTD GIGKSYAEEL AKHGMKVVLI SRSKDKLDQV SSEIKEKFKV ETRTIAVDFA SEDIYDKIKT GLAGLEIGIL VNNVGMSYEY PEYFLDVPDL DNVIKKMINI NILSVCKMTQ LVLPGMVERS KGAILNISSG SGMLPVPLLT IYSATKTFVD FFSQCLHEEY RSKGVFVQSV LPYFVATKLA KIRKPTLDKP SPETFVKSAI KTVGLQSRTN GYLIHALMGS IISNLPSWIY LKIVMNMNKS

TRAHYLKKTK KN

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human HSD17B12 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.	
Sterility:	0.22 μm filtered	
Endotoxin Level:	Protein is endotoxin-free.	
Grade:	Crystallography grade	

Target Details

Target: HSD17B12

Target Details

Alternative Name:	HSD17B12 (HSD17B12 Products)	
Background:	Catalyzes the second of the four reactions of the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process, allows the addition of two carbons to the chain of long- and very long-chain fatty acids/VLCFAs per cycle. This enzyme has a 3-ketoacyl-CoA reductase activity, reducing 3-ketoacyl-CoA to 3-hydroxyacyl-CoA, within each cycle of fatty acid elongation. Thereby, it may participate to the production of VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators. May also catalyze the transformation of estrone (E1) into estradiol (E2) and play a role in estrogen formation. {ECO:0000269 PubMed:12482854,	
	ECO:0000269 PubMed:16166196}.	
Molecular Weight:	35.5 kDa Including tag.	
UniProt:	Q53GQ0	
Pathways:	Steroid Hormone Biosynthesis	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process