

Datasheet for ABIN3115785

## CLEC4G Protein (AA 1-293) (Strep Tag)



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	1 mg
Target:	CLEC4G
Protein Characteristics:	AA 1-293
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CLEC4G protein is labelled with Strep Tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)

#### Product Details

Sequence: MDTTRYSKWG GSSEVPGGP WGRVHWSRR PLFLALAVLV TTVLWAVILS ILLSKASTER  
AALLDGHDLL RTNASKQTAA LGALKEEVGD CHSCCSGTQA QLQTTRAEELG EAQAKLMEQE  
SALRELRELV TQGLAEAGRG REDVRELFR ALEAVRLQNN SCEPCPTSWL SFEGSCYFFS  
VPKTTWAAAQ DHCADASAHL VIVGGLDEQG FLTRNTRGRG YWLGLRAVRH LGKVQGYQWV  
DGVSLFSHW NQGEPNDAWG RENCVMMLHT GLWNDAPCDS EKDGWICEKR HNC

**Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics: Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

---

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

---

#### Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

---

#### Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

---

## Product Details

---

Grade: Crystallography grade

## Target Details

---

Target: CLEC4G

Alternative Name: CLEC4G ([CLEC4G Products](#))

Background: C-type lectin domain family 4 member G (Liver and lymph node sinusoidal endothelial cell C-type lectin) (LSEctin),FUNCTION: Binds mannose, N-acetylglucosamine (GlcNAc) and fucose, but not galactose, in a Ca(2+)-dependent manner, in vitro. {ECO:0000269|PubMed:14711836}., FUNCTION: (Microbial infection) Acts as a receptor for Japanese encephalitis virus. {ECO:0000269|PubMed:24623090}., FUNCTION: (Microbial infection) Acts as a receptor for Ebolavirus. {ECO:0000269|PubMed:16051304}., FUNCTION: (Microbial infection) Acts as a receptor for SARS-CoV. {ECO:0000269|PubMed:16051304}., FUNCTION: (Microbial infection) Acts as a receptor for Lassa virus and Lymphocytic choriomeningitis virus glycoprotein (PubMed:22156524, PubMed:22673088). {ECO:0000269|PubMed:22156524, ECO:0000269|PubMed:22673088}.

Molecular Weight: 32.6 kDa

UniProt: [Q6UXB4](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

## Images

---



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process