antibodies

## Datasheet for ABIN3116214 STING/TMEM173 Protein (AA 1-379) (rho-1D4 tag)



Overview

2

Images

Quantity:	1 mg
Target:	STING/TMEM173 (TMEM173)
Protein Characteristics:	AA 1-379
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This STING/TMEM173 protein is labelled with rho-1D4 tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

## Product Details

Sequence:	MPHSSLHPSI PCPRGHGAQK AALVLLSACL VTLWGLGEPP EHTLRYLVLH LASLQLGLLL
	NGVCSLAEEL RHIHSRYRGS YWRTVRACLG CPLRRGALLL LSIYFYYSLP NAVGPPFTWM
	LALLGLSQAL NILLGLKGLA PAEISAVCEK GNFNVAHGLA WSYYIGYLRL ILPELQARIR
	TYNQHYNNLL RGAVSQRLYI LLPLDCGVPD NLSMADPNIR FLDKLPQQTG DHAGIKDRVY
	SNSIYELLEN GQRAGTCVLE YATPLQTLFA MSQYSQAGFS REDRLEQAKL FCRTLEDILA
	DAPESQNNCR LIAYQEPADD SSFSLSQEVL RHLRQEEKEE VTVGSLKTSA VPSTSTMSQE
	PELLISGMEK PLPLRTDFSG SSGTETSQVA PA
Specificity:	C-terminal Rho1D4-tag
Characteristics:	<ul> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> <li>Human TMEM173 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.</li> <li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li> </ul>

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3116214 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

## Product Details

	This made-to-order protein has already been successfully produced. Please let us know if you
	are interested in purchasing a smaller amount of this protein. We will check our stock and make
	you a customized quote in case we can provide this protein in a smaller amount
	When you order this made-to-order protein you will only pay upon receival of the correctly
	folded protein. With no financial risk on your end you can rest assured that our experienced
	protein experts will do everything to make sure that you receive the protein you ordered.
	The concentration of our recombinant proteins is measured using the absorbance at 280nm.
	The protein's absorbance will be measured in several dilutions and is measured against its
	specific reference buffer.
	The concentration of the protein is calculated using its specific absorption coefficient. We use
	the Expasy's protparam tool to determine the absorption coefficient of each protein.
Purification:	Three step purification of membrane proteins expressed in baculovirus infected SF9 insect
	cells:
	1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with
	different detergents (detergent screen). Samples are analyzed by Western blot.
	2. The best performing detergent is used for solubilization and the proteins are purified via their
	rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate
	fractions are analyzed by Western blot.
	3. Protein containing fractions of the best purification are subjected to second purification step
	through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and
	Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Protein is endotoxin-free.
Grade:	Crystallography grade
Target Details	
Target:	STING/TMEM173 (TMEM173)

l'alget.	0111(0), 11121117 0 (11121117 0)
Alternative Name:	TMEM173 (TMEM173 Products)
Background:	Facilitator of innate immune signaling that acts as a sensor of cytosolic DNA from bacteria and
	viruses and promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate
	immune response is triggered in response to non-CpG double-stranded DNA from viruses and
	bacteria delivered to the cytoplasm. Acts by recognizing and binding cyclic di-GMP (c-di-GMP),
	a second messenger produced by bacteria, and cyclic GMP-AMP (cGAMP), a messenger

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3116214 | 04/30/2024 | Copyright antibodies-online. All rights reserved.


	produced in response to DNA virus in the cytosol: upon binding of c-di-GMP or cGAMP,
	autoinhibition is alleviated and TMEM173/STING is able to activate both NF-kappa-B and IRF3
	transcription pathways to induce expression of type I interferon and exert a potent anti-viral
	state. May be involved in translocon function, the translocon possibly being able to influence
	the induction of type I interferons. May be involved in transduction of apoptotic signals via its
	association with the major histocompatibility complex class II (MHC-II). Mediates death
	signaling via activation of the extracellular signal-regulated kinase (ERK) pathway. Essential for
	the induction of IFN-beta in response to human herpes simplex virus 1 (HHV-1) infection.
	Exhibits 2',3' phosphodiester linkage-specific ligand recognition. Can bind both 2'-3' linked
	cGAMP and 3'-3' linked cGAMP but is preferentially activated by 2'-3' linked cGAMP
	(PubMed:26300263). {ECO:0000269 PubMed:18724357, ECO:0000269 PubMed:18818105,
	ECO:0000269 PubMed:19433799, ECO:0000269 PubMed:19776740,
	ECO:0000269 PubMed:21074459, ECO:0000269 PubMed:21947006,
	EC0:0000269 PubMed:23027953, EC0:0000269 PubMed:23258412,
	ECO:0000269 PubMed:23707065, ECO:0000269 PubMed:23722158,
	EC0:0000269 PubMed:26229117, EC0:0000269 PubMed:26300263}.
Molecular Weight:	43.4 kDa Including tag.
UniProt:	Q86WV6
Pathways:	Activation of Innate immune Response
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	20 mM Hepes, pH 8.0; 150 mM NaCl; 0,25 % LDAO + Protease inhibitors

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3116214 | 04/30/2024 | Copyright antibodies-online. All rights reserved.


Handlin	( 1
1 IGI IGIII I	ч

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

#### Images



TMEM173 (AA 1-379), gel filtration <u>Superose 6</u> in presence of protease inhibitors, fraction 12 and 13 identified with western blot



TMEM173 (AA 1-379), Fractions 12,13

# Size-exclusion chromatography-High Pressure Liquid Chromatography Image 1.

## Western Blotting

Image 2.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN3116214 | 04/30/2024 | Copyright antibodies-online. All rights reserved.