# antibodies .- online.com





# SLC13A5 Protein (AA 1-568) (Strep Tag)





Go to Product page

#### Overview

Quantity:	1 mg
Target:	SLC13A5
Protein Characteristics:	AA 1-568
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC13A5 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

#### **Product Details**

Sequence:

MASALSYVSK FKSFVILFVT PLLLLPLVIL MPAKFVRCAY VIILMAIYWC TEVIPLAVTS

LMPVLLFPLF QILDSRQVCV QYMKDTNMLF LGGLIVAVAV ERWNLHKRIA LRTLLWVGAK

PARLMLGFMG VTALLSMWIS NTATTAMMVP IVEAILQQME ATSAATEAGL ELVDKGKAKE

LPGSQVIFEG PTLGQQEDQE RKRLCKAMTL CICYAASIGG TATLTGTGPN VVLLGQMNEL

FPDSKDLVNF ASWFAFAFPN MLVMLLFAWL WLQFVYMRFN FKKSWGCGLE SKKNEKAALK

VLQEEYRKLG PLSFAEINVL ICFFLLVILW FSRDPGFMPG WLTVAWVEGE TKYVSDATVA

IFVATLLFIV PSQKPKFNFR SQTEEERKTP FYPPPLLDWK VTQEKVPWGI VLLLGGGFAL

AKGSEASGLS VWMGKQMEPL HAVPPAAITL ILSLLVAVFT ECTSNVATTT LFLPIFASMS

RSIGLNPLYI MLPCTLSASF AFMLPVATPP NAIVFTYGHL KVADMVKTGV IMNIIGVFCV

FLAVNTWGRA IFDLDHFPDW ANVTHIET

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

#### have a special request, please contact us.

#### Characteristics:

#### Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

## Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.

	<ol><li>Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li></ol>
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade
Target Details	
Target:	SLC13A5
Alternative Name:	SLC13A5 (SLC13A5 Products)
Background:	Na(+)/citrate cotransporter (NaCT) (Sodium-coupled citrate transporter) (Sodium-dependent
	citrate transporter) (Solute carrier family 13 member 5),FUNCTION: High-affinity sodium/citrate
	cotransporter that mediates the entry of citrate into cells, which is a critical participant of
	biochemical pathways (PubMed:12445824, PubMed:26324167, PubMed:26384929,
	PubMed:30054523, PubMed:33597751, PubMed:12826022). May function in various metabolic
	processes in which citrate has a critical role such as energy production (Krebs cycle), fatty acid
	synthesis, cholesterol synthesis, glycolysis, and gluconeogenesis (PubMed:12826022).
	Transports citrate into the cell in a Na(+)-dependent manner, recognizing the trivalent form of
	citrate (physiological pH ) rather than the divalent form (PubMed:12445824, PubMed:26324167
	PubMed:26384929, PubMed:30054523, PubMed:33597751, PubMed:12826022). Can recognize
	succinate as a substrate, but its affinity for succinate is several fold lower than for citrate
	(PubMed:26324167). The stoichiometry is probably 4 Na(+) for each carboxylate, irrespective or
	whether the translocated substrate is divalent or trivalent, rendering the process electrogenic
	(PubMed:12445824, PubMed:12826022). Involved in the regulation of citrate levels in the brain
	(By similarity). {ECO:0000250 UniProtKB:Q67BT3, ECO:0000269 PubMed:12445824,
	ECO:0000269 PubMed:12826022, ECO:0000269 PubMed:26324167,
	ECO:0000269 PubMed:26384929, ECO:0000269 PubMed:30054523,
	ECO:0000269 PubMed:33597751}.
Molecular Weight:	63.1 kDa
UniProt:	Q86YT5
Pathways:	Dicarboxylic Acid Transport

## **Application Details**

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
	During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process