antibodies .- online.com





UNC5B Protein (AA 27-945) (rho-1D4 tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	UNC5B
Protein Characteristics:	AA 27-945
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This UNC5B protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

Sequence:

GTDSGSEVLP DSFPSAPAEP LPYFLQEPQD AYIVKNKPVE LRCRAFPATQ IYFKCNGEWV SQNDHVTQEG LDEATGLRVR EVQIEVSRQQ VEELFGLEDY WCQCVAWSSA GTTKSRRAYV RIAYLRKNFD QEPLGKEVPL DHEVLLQCRP PEGVPVAEVE WLKNEDVIDP TQDTNFLLTI DHNLIIRQAR LSDTANYTCV AKNIVAKRRS TTATVIVYVN GGWSSWAEWS PCSNRCGRGW QKRTRTCTNP APLNGGAFCE GQAFQKTACT TICPVDGAWT EWSKWSACST ECAHWRSREC MAPPPQNGGR DCSGTLLDSK NCTDGLCMQN KKTLSDPNSH LLEASGDAAL YAGLVVAIFV VVAILMAVGV VVYRRNCRDF DTDITDSSAA LTGGFHPVNF KTARPSNPQL LHPSVPPDLT ASAGIYRGPV YALQDSTDKI PMTNSPLLDP LPSLKVKVYS SSTTGSGPGL ADGADLLGVL PPGTYPSDFA RDTHFLHLRS ASLGSQQLLG LPRDPGSSVS GTFGCLGGRL SIPGTGVSLL VPNGAIPQGK FYEMYLLINK AESTLPLSEG TQTVLSPSVT CGPTGLLLCR PVILTMPHCA EVSARDWIFQ LKTQAHQGHW EEVVTLDEET LNTPCYCQLE PRACHILLDQ LGTYVFTGES YSRSAVKRLQ LAVFAPALCT SLEYSLRVYC LEDTPVALKE VLELERTLGG YLVEEPKPLM

FKDSYHNLRL SLHDLPHAHW RSKLLAKYQE IPFYHIWSGS QKALHCTFTL ERHSLASTEL
TCKICVRQVE GEGQIFQLHT TLAETPAGSL DTLCSAPGST VTTQLGPYAF KIPLSIRQKI
CNSLDAPNSR GNDWRMLAQK LSMDRYLNYF ATKASPTGVI LDLWEALQQD DGDLNSLASA
LEEMGKSEML VAVATDGDC

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human UNC5B Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- 3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and

Product Details

ed by SDS PAGE, Size Exclusion Chromatography and Western Blot. In-free. ade roducts) In required for axon guidance. Mediates axon repulsion of neuronal growth
n-free. ade
ade roducts)
ade roducts)
roducts)
<u> </u>
<u> </u>
<u> </u>
required for axon guidance. Mediates axon repulsion of neuronal growth
oping nervous system upon ligand binding. Axon repulsion in growth cones its association with DCC that may trigger signaling for repulsion (By ns as netrin receptor that negatively regulates vascular branching during liates retraction of tip cell filopodia on endothelial growth cones in response rity). It also acts as a dependence receptor required for apoptosis induction ed with netrin ligand (PubMed:12598906). Mediates apoptosis by activating ence of NTN1, activates DAPK1 by reducing its autoinhibitory: Ser-308 thereby increasing its catalytic activity (By similarity). ProtKB:008722, ECO:0000250 UniProtKB:Q8K1S3, Med:12598906}.
g tag.
pplications listed above we expect the protein to work for functional studies tein has not been tested for functional studies yet we cannot offer a gurantee

Application Details

Expiry Date:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Unlimited (if stored properly)