
antibodies .- online.com





TRPM4 Protein (AA 1-1214) (rho-1D4 tag)





Go to Product page

Overview

Quantity:	1 mg
Target:	TRPM4
Protein Characteristics:	AA 1-1214
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRPM4 protein is labelled with rho-1D4 tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), Crystallization (Crys), ELISA

Product Details

Sequence:

MVVPEKEQSW IPKIFKKKTC TTFIVDSTDP GGTLCQCGRP RTAHPAVAME DAFGAAVVTV
WDSDAHTTEK PTDAYGELDF TGAGRKHSNF LRLSDRTDPA AVYSLVTRTW GFRAPNLVVS
VLGGSGGPVL QTWLQDLLRR GLVRAAQSTG AWIVTGGLHT GIGRHVGVAV RDHQMASTGG
TKVVAMGVAP WGVVRNRDTL INPKGSFPAR YRWRGDPEDG VQFPLDYNYS AFFLVDDGTH
GCLGGENRFR LRLESYISQQ KTGVGGTGID IPVLLLLIDG DEKMLTRIEN ATQAQLPCLL
VAGSGGAADC LAETLEDTLA PGSGGARQGE ARDRIRRFFP KGDLEVLQAQ VERIMTRKEL
LTVYSSEDGS EEFETIVLKA LVKACGSSEA SAYLDELRLA VAWNRVDIAQ SELFRGDIQW
RSFHLEASLM DALLNDRPEF VRLLISHGLS LGHFLTPMRL AQLYSAAPSN SLIRNLLDQA
SHSAGTKAPA LKGGAAELRP PDVGHVLRML LGKMCAPRYP SGGAWDPHPG QGFGESMYLL
SDKATSPLSL DAGLGQAPWS DLLLWALLLN RAQMAMYFWE MGSNAVSSAL GACLLLRVMA
RLEPDAEEAA RRKDLAFKFE GMGVDLFGEC YRSSEVRAAR LLLRRCPLWG DATCLQLAMQ
ADARAFFAQD GVQSLLTQKW WGDMASTTPI WALVLAFFCP PLIYTRLITF RKSEEEPTRE

ELEFDMDSVI NGEGPVGTAD PAEKTPLGVP RQSGRPGCCG GRCGGRRCLR RWFHFWGAPV
TIFMGNVVSY LLFLLLFSRV LLVDFQPAPP GSLELLLYFW AFTLLCEELR QGLSGGGGSL
ASGGPGPGHA SLSQRLRLYL ADSWNQCDLV ALTCFLLGVG CRLTPGLYHL GRTVLCIDFM
VFTVRLLHIF TVNKQLGPKI VIVSKMMKDV FFFLFFLGVW LVAYGVATEG LLRPRDSDFP
SILRRVFYRP YLQIFGQIPQ EDMDVALMEH SNCSSEPGFW AHPPGAQAGT CVSQYANWLV
VLLLVIFLLV ANILLVNLLI AMFSYTFGKV QGNSDLYWKA QRYRLIREFH SRPALAPPFI
VISHLRLLLR QLCRRPRSPQ PSSPALEHFR VYLSKEAERK LLTWESVHKE NFLLARARDK
RESDSERLKR TSQKVDLALK QLGHIREYEQ RLKVLEREVQ QCSRVLGWVA EALSRSALLP
PGGPPPPDLP GSKD

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human TRPM4 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with

- different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- 3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

 Purity:
 >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

 Sterility:
 0.22 μm filtered

 Endotoxin Level:
 Protein is endotoxin-free.

 Grade:
 Crystallography grade

Target Details

Target:	TRPM4			
Alternative Name:	TRPM4 (TRPM4 Products)			
Background:	Calcium-activated non selective (CAN) cation channel that mediates membrane depolarization			
	While it is activated by increase in intracellular Ca(2+), it is impermeable to it. Mediates			
	transport of monovalent cations $(Na(+) > K(+) > Cs(+) > Li(+))$, leading to depolarize the			
	membrane. It thereby plays a central role in cadiomyocytes, neurons from entorhinal cortex,			
	dorsal root and vomeronasal neurons, endocrine pancreas cells, kidney epithelial cells, cochlea			
	hair cells etc. Participates in T-cell activation by modulating Ca(2+) oscillations after T			
	lymphocyte activation, which is required for NFAT-dependent IL2 production. Involved in			
	myogenic constriction of cerebral arteries. Controls insulin secretion in pancreatic beta-cells.			
	May also be involved in pacemaking or could cause irregular electrical activity under conditions			
	of Ca(2+) overload. Affects T-helper 1 (Th1) and T-helper 2 (Th2) cell motility and cytokine			
	production through differential regulation of calcium signaling and NFATC1 localization.			
	Enhances cell proliferation through up-regulation of the beta-catenin signaling pathway.			
	{ECO:0000269 PubMed:12015988, ECO:0000269 PubMed:12799367,			
	ECO:0000269 PubMed:15121803, ECO:0000269 PubMed:15472118,			
	ECO:0000269 PubMed:15550671, ECO:0000269 PubMed:16806463,			
	ECO:0000269 PubMed:20625999, ECO:0000269 PubMed:20656926}.			
Molecular Weight:	135.5 kDa Including tag.			
UniProt:	Q8TD43			

Target Details

\mathbf{P}	at	n	۱۱۸	12	١/	c.

Regulation of Leukocyte Mediated Immunity, Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process