

Datasheet for ABIN3117748
NDFIP1 Protein (AA 2-221) (rho-1D4 tag)[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	NDFIP1
Protein Characteristics:	AA 2-221
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NDFIP1 protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

Sequence:	<p>ALALAALAAV EPACGSRYQQ LQNEEESGEP EQAAGDAPPP YSSISAESAA YFDYKDESGF PKPPSYNVAT TLPSYDEAER TKAEATIPLV PGRDEDFVGR DDFDDADQLR IGNDGIFMLT FFMAFLFNWI GFFLSFCLTT SAAGRYGAIS GFGLSLIKWI LIVRFSTYFP GYFDGQYWLLW WVFLVLGFL FLRGFINYAK VRKMPETFSN LPRTRVLFYI</p> <p>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</p>
Characteristics:	<ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Human NDFIP1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.• State-of-the-art algorithm used for plasmid design (Gene synthesis). <p>This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.</p>

Product Details

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's protParam tool to determine the absorption coefficient of each protein.

Purification:	Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none">1. Membrane proteins are fractionated by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
---------------	--

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
---------	--

Sterility:	0.22 µm filtered
------------	------------------

Endotoxin Level:	Protein is endotoxin-free.
------------------	----------------------------

Grade:	Crystallography grade
--------	-----------------------

Target Details

Target:	NDFIP1
Alternative Name:	NDFIP1 (NDFIP1 Products)
Background:	Activates HECT domain-containing E3 ubiquitin-protein ligases, including NEDD4 and ITCH, and

Target Details

consequently modulates the stability of their targets. As a result, controls many cellular processes. Prevents chronic T-helper cells-mediated inflammation by activating ITCH and thus controlling JUNB degradation (By similarity). In cortical neurons, mediates the ubiquitination of SLC11A2/DMT1 by NEDD4L, leading to down-regulation of the divalent metal transporter and protection of the cells from cobalt and iron toxicity (PubMed:19706893). Modulates EGFR signaling through multiple pathways. In particular, may regulate the ratio of AKT1-to-MAPK8 signaling in response to EGF, acting on AKT1 probably through PTEN destabilization and on MAPK8 through ITCH-dependent MAP2K4 inactivation. As a result, may control cell growth rate (PubMed:20534535). Enhances the ubiquitination of BRAT1 mediated by E3 ubiquitin-protein ligases: NEDD4, NEDD4L and ITCH, and is required for the nuclear localization of ubiquitinated BRAT1 (PubMed:25631046). {ECO:0000250|UniProtKB:Q8R0W6, ECO:0000269|PubMed:19343052, ECO:0000269|PubMed:19706893, ECO:0000269|PubMed:20534535, ECO:0000269|PubMed:25631046}.

Molecular Weight: 25.9 kDa Including tag.

UniProt: [Q9BT67](#)

Pathways: [Transition Metal Ion Homeostasis](#), [Regulation of Leukocyte Mediated Immunity](#), [Production of Molecular Mediator of Immune Response](#), [Negative Regulation of Transporter Activity](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Handling

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process