

Datasheet for ABIN3117971

Der1-Like Domain Family, Member 2 (DERL2) (AA 1-239) protein (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	Der1-Like Domain Family, Member 2 (DERL2)
Protein Characteristics:	AA 1-239
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	Strep Tag
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Brand:	ALiCE®
Sequence:	<p>MAYQSLRLEY LQIPVSRAY TTACVLTTAA VQLELITPFQ LYFNPELIFK HFQIWRLITN FLFFGPVGFN FLNMFILYR YCRMLEEGSF RGRTADVFVM FLFGGFLMTL FGLFVSLVFL GQAFITIMLVY VWSRRNPYVR MNFFGLLNFQ APFLPWVLMG FSLLLGNSII VDLLGIAVGH IYFFLEDVFP NQPGGIRILK TPSILKAIFD TPDEDPNYPN LPEERPGGFA WGEQGRLGG</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"> • Made in Germany - from design to production - by highly experienced protein experts. • Protein expressed with ALiCE® and purified in one-step affinity chromatography • These proteins are normally active (enzymatically functional) as our customers have

- reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	Der1-Like Domain Family, Member 2 (DERL2)
Alternative Name:	DERL2 (DERL2 Products)
Background:	Derlin-2 (Degradation in endoplasmic reticulum protein 2) (DERtrin-2) (Der1-like protein 2) (F-

Target Details

LAN-1) (F-LANa),FUNCTION: Functional component of endoplasmic reticulum-associated degradation (ERAD) for misfolded luminal glycoproteins, but not that of misfolded nonglycoproteins. May act by forming a channel that allows the retrotranslocation of misfolded glycoproteins into the cytosol where they are ubiquitinated and degraded by the proteasome. May mediate the interaction between VCP and misfolded glycoproteins (PubMed:16186509, PubMed:16449189). May also be involved in endoplasmic reticulum stress-induced pre-emptive quality control, a mechanism that selectively attenuates the translocation of newly synthesized proteins into the endoplasmic reticulum and reroutes them to the cytosol for proteasomal degradation (PubMed:26565908). {ECO:0000269|PubMed:16186509, ECO:0000269|PubMed:16449189, ECO:0000269|PubMed:26565908}., FUNCTION: (Microbial infection) In contrast to DERL1, it is not involved in the degradation of MHC class I heavy chains following infection by cytomegaloviruses. {ECO:0000269|PubMed:15215855}.

Molecular Weight: 27.6 kDa

UniProt: [Q9GZP9](#)

Pathways: [ER-Nucleus Signaling, Feeding Behaviour](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months