antibodies .- online.com





CD57 Protein (AA 1-334) (Strep Tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	CD57 (B3GAT1)
Protein Characteristics:	AA 1-334
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD57 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)

Product Details

Sequence:

MPKRRDILAI VLIVLPWTLL ITVWHQSTLA PLLAVHKDEG SDPRRETPPG ADPREYCTSD RDIVEVVRTE YVYTRPPPWS DTLPTIHVVT PTYSRPVQKA ELTRMANTLL HVPNLHWLVV EDAPRRTPLT ARLLRDTGLN YTHLHVETPR NYKLRGDARD PRIPRGTMQR NLALRWLRET FPRNSSQPGV VYFADDDNTY SLELFEEMRS TRRVSVWPVA FVGGLRYEAP RVNGAGKVVG WKTVFDPHRP FAIDMAGFAV NLRLILQRSQ AYFKLRGVKG GYQESSLLRE LVTLNDLEPK AANCTKILVW HTRTEKPVLV NEGKKGFTDP SVEI

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

correct folding and modification.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details	
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade
Target Details	
Target:	CD57 (B3GAT1)
Alternative Name:	B3GAT1 (B3GAT1 Products)
Background:	Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1 (EC 2.4.1.135) (Beta-1,3-glucuronyltransferase 1) (Glucuronosyltransferase P) (GlcAT-P) (UDP-GlcUA:glycoprotein beta-1,3-glucuronyltransferase) (GlcUAT-P),FUNCTION: Involved in the biosynthesis of L2/HNK-1 carbohydrate epitope on glycoproteins. Can also play a role in glycosaminoglycan biosynthesis. Substrates include asialo-orosomucoid (ASOR), asialo-fetuin, and asialo-neural cell adhesion molecule. Requires sphingomyelin for activity: stearoyl-sphingomyelin was the most effective, followed by palmitoyl-sphingomyelin and lignoceroyl-sphingomyelin. Activity was demonstrated only for sphingomyelin with a saturated fatty acid and not for that with an unsaturated fatty acid, regardless of the length of the acyl group. {ECO:0000250 UniProtKB:035789}.
Molecular Weight:	38.3 kDa
UniProt:	Q9P2W7
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional

components needed for protein production (amino acids, cofactors, etc.) are added to produce

something that functions like a cell, but without the constraints of a living system - all that's

Application Details

	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)
Images	




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process