antibodies .- online.com



Datasheet for ABIN3119310

Chromosome 11open Reading Frame 9 (C11orf9) (AA 587-1151) protein (rho-1D4 tag)



Go to Product page



Overview

Quantity:	1 mg
Target:	Chromosome 11open Reading Frame 9 (C11orf9)
Protein Characteristics:	AA 587-1151
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	rho-1D4 tag
Application:	Crystallization (Crys), ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:

SDLRAKEHVQ EVDTTEQLKR ISRMRLVHYR YKPEFAASAG IEATAPETGV IAQEVKEILP
EAVKDTGDMV FANGKTIENF LVVNKERIFM ENVGAVKELC KLTDNLETRI DELERWSHKL
AKLRRLDSLK STGSSGAFSH AGSQFSRAGS VPHKKRPPKV ASKSSSVVPD QACISQRFLQ
GTIIALVVVM AFSVVSMSTL YVLSLRTEED LVDTDGSFAV STSCLLALLR PQPPGGSEAL
CPWSSQSFGT TQLRQSPLTT GLPGIQPSLL LVTTSLTSSA PGSAVRTLDM CSSHPCPVIC
CSSPTTNPTT GPSLGPSFNP GHVLSPSPSP STNRSGPSQM ALLPVTNIRA KSWGLSVNGI
GHSKHHKSLE PLASPAVPFP GGQGKAKNSP SLGFHGRARR GALQSSVGPA EPTWAQGQSA
SLLAEPVPSL TSIQVLENSM SITSQYCAPG DACRPGNFTY HIPVSSGTPL HLSLTLQMNS
SSPVSVVLCS LRSKEEPCEE GSLPQSLHTH QDTQGTSHRW PITILSFREF TYHFRVALLG
QANCSSEALA QPATDYHFHF YRLCD

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human MYRF Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- 3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

 Purity:
 >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

 Sterility:
 0.22 μm filtered

 Endotoxin Level:
 Protein is endotoxin-free.

 Grade:
 Crystallography grade

Target Details

Target:	Chromosome 11open Reading Frame 9 (C11orf9)
Alternative Name:	MYRF (C11orf9 Products)
Background:	Myelin regulatory factor: Consitutes a precursor of the transcription factor. Mediates the
	autocatalytic cleavage that releases the Myelin regulatory factor, N-terminal component that
	specifically activates transcription of central nervous system (CNS) myelin genes
	(PubMed:23966832). {ECO:0000269 PubMed:23966832}., Myelin regulatory factor, C-terminal:
	Membrane-bound part that has no transcription factor activity and remains attached to the
	endoplasmic reticulum membrane following cleavage. {ECO:0000269 PubMed:23966832}.,
	Myelin regulatory factor, N-terminal: Transcription factor that specifically activates expression
	of myelin genes such as MBP, MOG, MAG and PLP1 during oligodendrocyte (OL) maturation,
	thereby playing a central role in oligodendrocyte maturation and CNS myelination. Specifically
	recognizes and binds DNA sequence 5'-CTGGYAC-3' in the regulatory regions of myelin-specific
	genes and directly activates their expression. Not only required during oligodendrocyte
	differentiation but is also required on an ongoing basis for the maintenance of expression of
	myelin genes and for the maintenance of a mature, viable oligodendrocyte phenotype
	(PubMed:23966832). {ECO:0000269 PubMed:23966832}.
Molecular Weight:	61.8 kDa Including tag.
UniProt:	Q9Y2G1
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
	For Research Use only
Restrictions:	
Restrictions: Handling	
	Liquid

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process