# antibodies -online.com





## BCL2L10 Protein (AA 1-191) (Strep Tag)



Go to Product page

( )	11/0	K\ /	iew
	$\cup$	ועוי	$I \cap VV$

Quantity:	1 mg
Target:	BCL2L10
Protein Characteristics:	AA 1-191
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This BCL2L10 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB)

#### **Product Details**

#### Sequence:

MADSQDPLHE RTRRLLSDYI FFCAREPDTP EPPPTSVEAA LLRSVTRQIQ QEHQEFFSSF
CESRGNRLEL VKQMADKLLS KDQDFSWSQL VMLLAFAGTL MNQGPYMAVK QKRDLGNRVI
VTRDCCLIVN FLYNLLMGRR HRARLEALGG WDGFCRFFKN PLPLGFWRRL LIQAFLSGFF
ATAIFFIWKR L

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- · These proteins are normally active (enzymatically functional) as our customers have

reported (not tested by us and not guaranteed).

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### **Expression System:**

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

### Target Details

Alternative Name: Background:		
Background:	Bcl2l10 (BCL2L10 Products)	
	Bcl-2-like protein 10 (Bcl2-L-10) (Anti-apoptotic protein Boo) (Apoptosis regulator Bcl-B) (Bcl-2 homolog Diva),FUNCTION: Promotes cell survival by suppressing apoptosis induced by BAX but not BAK (By similarity). Increases binding of AHCYL1/IRBIT to ITPR1 (By similarity). Reduces ITPR1-mediated calcium release from the endoplasmic reticulum cooperatively with AHCYL1/IRBIT under normal cellular conditions (By similarity). Under apoptotic stress conditions, dissociates from ITPR1 and is displaced from mitochondria-associated endoplasmic reticulum membranes, leading to increased Ca(2+) transfer to mitochondria which promotes apoptosis (By similarity). Required for the correct formation of the microtubule organizing center during oocyte cell division, potentially via regulation of protein abundance and localization of other microtubule organizing center components such as AURKA and TPX2 (PubMed:27753540). {ECO:0000250 UniProtKB:Q9HD36, ECO:0000269 PubMed:27753540}.	
Molecular Weight:	22.3 kDa	
UniProt:	Q9Z0F3	
Pathways:  Application Details	Positive Regulation of Endopeptidase Activity	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the	
	mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!	

## Handling

Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	