

Datasheet for ABIN3120404
TSC22D3 Protein (AA 1-137) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	TSC22D3
Protein Characteristics:	AA 1-137
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TSC22D3 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence: MNTEMYQTPM EVAVYQLHNF SISFFSSLLG GDVVSVKLDN SASGASVVAL DNKIEQAMD
L VKNHLMYAVR EEVEVLKEQI RELLEKNSQL ERENTLLKTL ASPEQLEKFQ SRLSPEEPAP
EAPETPETPE APGGSVA

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Protein expressed with ALiCE® and purified in one-step affinity chromatography• These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).• State-of-the-art algorithm used for plasmid design (Gene synthesis).
------------------	---

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
---------------	--

Purity:	> 80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
---------	---

Target Details

Target:	TSC22D3
---------	---------

Alternative Name:	Tsc22d3 (TSC22D3 Products)
-------------------	--

Background:	TSC22 domain family protein 3 (Glucocorticoid-induced leucine zipper protein) (TSC22-related-inducible leucine zipper 3) (Tilz3),FUNCTION: Protects T-cells from IL2 deprivation-induced apoptosis through the inhibition of FOXO3A transcriptional activity that leads to the down-regulation of the pro-apoptotic factor BCL2L11 (PubMed:26752201). In macrophages, plays a
-------------	---

Target Details

role in the anti-inflammatory and immunosuppressive effects of glucocorticoids and IL10 (By similarity). In T-cells, inhibits anti-CD3-induced NFKB1 nuclear translocation and thereby NFKB1 DNA-binding activities (By similarity). In vitro, suppresses AP-1 transcription factor complex DNA-binding activities (PubMed:11397794). {ECO:0000250|UniProtKB:Q99576, ECO:0000269|PubMed:11397794, ECO:0000269|PubMed:26752201}., FUNCTION: [Isoform 1]: Inhibits myogenic differentiation and mediates anti-myogenic effects of glucocorticoids by binding and regulating MYOD1 and HDAC1 transcriptional activity resulting in reduced expression of MYOG. {ECO:0000269|PubMed:20124407}., FUNCTION: [Isoform 4]: Inhibits myogenic differentiation and mediates anti-myogenic effects of glucocorticoids by binding and regulating MYOD1 and HDAC1 transcriptional activity resulting in reduced expression of MYOG. {ECO:0000269|PubMed:20124407}.

Molecular Weight: 15.2 kDa

UniProt: [Q9Z2S7](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer. If you have a special request,

Handling

please contact us.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)