

Datasheet for ABIN3120566

B3GAT3 Protein (AA 1-335) (Strep Tag)



Overview

Quantity:	250 μg
Target:	B3GAT3
Protein Characteristics:	AA 1-335
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This B3GAT3 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details	
Brand:	AliCE®
Sequence:	MKLKLKNVFL AYFLVSIAGL LYALVQLGQP CDCLPPLRAA AEQLRQKDLR ISQLQADLRR
	PPPVPAQPPE PEALPTIYVI TPTYARLVQK AELVRLSQTL SLVPRLHWLL VEDAESPTPL
	VSGLLAASGL LFTHLAVLTP KAQRLREGEP GWVRPRGVEQ RNKALDWLRG KGGAVGGEKD
	PPPPGTQGVV YFADDDNTYS RELFKEMRWT RGVSVWPVGL VGGLRFEGPQ VQDGRVVGFH
	TAWEPNRPFP LDMAGFAVAL PLLLAKPNAQ FDATAPRGHL ESSLLSHLVD PKDLEPRAAN
	CTQVLVWHTR TEKPKMKQEE QLQRQGQGSD PAIEV
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	B3GAT3

Target Details

Alternative Name:	B3gat3 (B3GAT3 Products)
Background:	Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 3 (EC 2.4.1.135) (Beta-1,3-
	glucuronyltransferase 3) (Glucuronosyltransferase I) (GlcAT-I) (UDP-GlcUA:Gal beta-1,3-Gal-R
	glucuronyltransferase) (GlcUAT-I),FUNCTION: Glycosaminoglycans biosynthesis. Involved in
	forming the linkage tetrasaccharide present in heparan sulfate and chondroitin sulfate.
	Transfers a glucuronic acid moiety from the uridine diphosphate-glucuronic acid (UDP-GlcUA)
	to the common linkage region trisaccharide Gal-beta-1,3-Gal-beta-1,4-Xyl covalently bound to a
	Ser residue at the glycosaminylglycan attachment site of proteoglycans. Can also play a role in
	the biosynthesis of I2/HNK-1 carbohydrate epitope on glycoproteins. Stimulates 2-
	phosphoxylose phosphatase activity of PXYLP1 in presence of uridine diphosphate-glucuronic
	acid (UDP-GlcUA) during completion of linkage region formation.
	{ECO:0000250 UniProtKB:094766}.
Molecular Weight:	37.1 kDa
JniProt:	P58158
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Dantuintinun	For Decearch Lieu only
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months