antibodies .- online.com

Datasheet for ABIN3120884 TUBA1C Protein (AA 1-449) (His tag)

1 Image



Overview

Quantity:	1 mg
Target:	TUBA1C
Protein Characteristics:	AA 1-449
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TUBA1C protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

TVVPGGDLAK VQRAVCMLSN TTAIAEAWAR LDHKFDLM AREDMAALEK DYEEVGADSA EGDDEGEEY
Sequence without tag. Tag location is at the discretion
special request, please contact us.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3120884 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you ander this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:1. In a first purification step, the protein is purified from the cleared cell ysate using three different His-tag capture materials: high yield. ED1A resistant, Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Fluste fractions are analyzed by SDS-PAGE and Western blot.Purification:> 295 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. <th></th> <th></th>		
The big advantage of ordering our made-to-order proteins in companison to ordering oustom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experimened protein scatter will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Express's protearam tool to determine the absorption coefficient of each protein. Purification: Two step purification of proteins expressed in baculovirus infected SP9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysale using three different His tage capture materials. high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PACE and Western Blot. Purify: >95 % as determined by SDS PAGE. 2. Protein containing fractions of the best purification are subjected to second purificatio		This protein is a made to order protein and will be made for the first time for your order. Our
made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Etuate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE 2. Protein containing fractions of the best purification are analyzed by SDS-PAGE 3. Purity:Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Target DetailsTurget DetailsTarget DetailsTUBA1C		experts in the lab will ensure that you receive a correctly folded protein.
cannot be expressed or purified.In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification step, the protein is purified from the cleared cell lysate using three different His tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PACE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purify.>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		The big advantage of ordering our made-to-order proteins in comparison to ordering custom
In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBATC		made proteins from other companies is that there is no financial obligation in case the protein
(other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials. high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purify:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		cannot be expressed or purified.
Purification: Usedep proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expassy's protparam tool to determine the absorption coefficient of each protein. Purification: Two step purification step, the protein is purified from the cleared cell lysate using three different Histag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade Target Details TUBA1C		In the unlikely event that the protein cannot be expressed or purified we do not charge anything
experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different Histag capture materials. high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTuBA1C		(other companies might charge you for any performed steps in the expression process for
When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different Histag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression
folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget:TUBA1C		experiments or purification optimization).
protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterillty:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		When you order this made-to-order protein you will only pay upon receival of the correctly
The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		folded protein. With no financial risk on your end you can rest assured that our experienced
The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		protein experts will do everything to make sure that you receive the protein you ordered.
specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		The concentration of our recombinant proteins is measured using the absorbance at 280nm.
The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.Purify:-Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:-95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		The protein's absorbance will be measured in several dilutions and is measured against its
the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		specific reference buffer.
Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		The concentration of the protein is calculated using its specific absorption coefficient. We use
1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		the Expasy's protparam tool to determine the absorption coefficient of each protein.
different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C	Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		1. In a first purification step, the protein is purified from the cleared cell lysate using three
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		
through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTuBA1C		
Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTUBA1C		
Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade Target Details TuBA1C		Western blot.
Endotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget DetailsTuBA1C	Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Grade: Crystallography grade Target Details Tuba1C	Sterility:	0.22 µm filtered
Target Details Target: TUBA1C	Endotoxin Level:	Protein is endotoxin free.
Target: TUBA1C	Grade:	Crystallography grade
Target: TUBA1C	Target Details	
		ΤΠΒΦΙΟ
Alternative Name: Tuba1c (TUBA1C Products)		
	Alternative Name:	Tuba1c (TUBA1C Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3120884 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.
Molecular Weight:	50.9 kDa Including tag.
UniProt:	P68373
Pathways:	Microtubule Dynamics, M Phase
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3120884 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN3120884 | 09/11/2023 | Copyright antibodies-online. All rights reserved.