

Datasheet for ABIN3120908

PIP5K1B Protein (AA 1-539) (Strep Tag)



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	250 μg
Target:	PIP5K1B
Protein Characteristics:	AA 1-539
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PIP5K1B protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details	
Brand:	AliCE®
Sequence:	MSSTAENGDA VPGKQNEEKT YKKTASSAIK GAIQLGIGYT VGNLTSKPER DVLMQDFYVV
	ESVFLPSEGS NLTPAHHYPD FRFKTYAPLA FRYFRELFGI KPDDYLYSIC SEPLIELSNP
	GASGSLFFLT SDDEFIIKTV QHKEAEFLQK LLPGYYMNLN QNPRTLLPKF YGLYCMQSGG
	INIRIVVMNN VLPRAMRMHL TYDLKGSTYK RRASRKEREK PNPTFKDLDF LQDMHEGLYF
	DTETYNALMK TLQRDCRVLE SFKIMDYSLL LGIHILDHSL KDKEEEPLQN VPDAKRPGMQ
	KVLYSTAMES IQGPGKSADG IIAENPDTMG GIPAKSHKGE KLLLFMGIID ILQSYRLMKK
	LEHSWKALVY DGDTVSVHRP SFYADRFLKF MNSRVFKKIQ ALKASPSKKR CNSIAALKAT
	SQEIVSSISQ EWKDEKRDLL TEGQSFSSLD EEALGSRHRP DLVPSTPSLF EAASLATTIS
	SSSLYVGEHY PHDRTTLYSN SKGLPSSSTF TLEEGTIYLT AEPNTLDLQD DASVLDVYL
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	PIP5K1B		
Alternative Name:	Pip5k1b (PIP5K1B Products)		
Background:	Phosphatidylinositol 4-phosphate 5-kinase type-1 beta (PIP5KI-beta) (PtdIns(4)P-5-kinase 1		
	beta) (EC 2.7.1.68) (68 kDa type I phosphatidylinositol 4-phosphate 5-kinase beta)		
	(Phosphatidylinositol 4-phosphate 5-kinase type I alpha) (PIP5KIalpha) (Phosphatidylinositol 4-		
	phosphate 5-kinase type I beta) (PIP5KIbeta),FUNCTION: Catalyzes the phosphorylation of		
	phosphatidylinositol 4-phosphate (PtdIns(4)P/PI4P) to form phosphatidylinositol 4,5-		
	bisphosphate (PtdIns(4,5)P2/PIP2), a lipid second messenger that regulates several cellular		
	processes such as signal transduction, vesicle trafficking, actin cytoskeleton dynamics, cell		
	adhesion, and cell motility (PubMed:8798574, PubMed:9367159, PubMed:9535851,		
	PubMed:22942276). PtdIns(4,5)P2 can directly act as a second messenger or can be utilized as		
	a precursor to generate other second messengers: inositol 1,4,5-trisphosphate (IP3),		
	diacylglycerol (DAG) or phosphatidylinositol-3,4,5-trisphosphate (PtdIns(3,4,5)P3/PIP3) (By		
	similarity). Mediates RAC1-dependent reorganization of actin filaments (PubMed:10679324).		
	Contributes to the activation of phospholipase PLD2 (PubMed:11032811). Together with		
	PIP5K1A, is required, after stimulation by G-protein coupled receptors, for the synthesis of IP3		
	that will induce stable platelet adhesion (PubMed:18772378).		
	{ECO:0000250 UniProtKB:Q99755, ECO:0000269 PubMed:10679324,		
	ECO:0000269 PubMed:11032811, ECO:0000269 PubMed:18772378,		
	ECO:0000269 PubMed:22942276, ECO:0000269 PubMed:8798574,		
	ECO:0000269 PubMed:9367159, ECO:0000269 PubMed:9535851}.		
Molecular Weight:	60.8 kDa		
UniProt:	P70181		
Pathways:	PI3K-Akt Signaling, Inositol Metabolic Process, Cell-Cell Junction Organization		
Application Details			
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies		
	as well. As the protein has not been tested for functional studies yet we cannot offer a		
	guarantee though.		
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from		
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce		
	even the most difficult-to-express proteins, including those that require post-translational		
	modifications.		

Application Details

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months