

Datasheet for ABIN3120974

TANK Protein (AA 1-448) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	TANK
Protein Characteristics:	AA 1-448
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TANK protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MSLKRHSLRR NACHLETRAG IPTILYSDAT GQRGMDKNIG EQLNRAYEAF RQACMDRDSA VRELQKQKTE NYEQRIREQQ EQLSFQQNLI DRLKSQLLL V DSSRDNSYGY VPLLEDSDRR KNNLTLDPEH DKVKLGTLRD KQSKVRRQEV SSGKESAKGL NIPLHHERDN IEKTFWDLKE EFHRICLLAK AQKDHL SKLN IPDIATDTQC SVPIQCTDKT EKQEALFKPQ AKDDINRGMS CVTAVTPRGL GRDEEDTSFE SLSKFNVKFP PMDNDSIFLH STPEAPSILA PATPETVCQD RFNMEVRDNP GNFVKTEETL FEIQGIDPIT SAIQNLKTTD KTNPSNL RAT CLPAGDHN VF YVNTFPLQDP PDAPFSLDS PGKAVRGPQQ PFWKPFLNQD TDLVVPDSD SELLKPLVCE FCQELFPPSI TSRGDFLRHL NTHFNGET</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>

Product Details

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	TANK
Alternative Name:	Tank (TANK Products)
Background:	<p>TRAF family member-associated NF-kappa-B activator (TRAF-interacting protein) (I-TRAF),FUNCTION: Adapter protein involved in I-kappa-B-kinase (IKK) regulation which constitutively binds TBK1 and IKBKE playing a role in antiviral innate immunity. Acts as a regulator of TRAF function by maintaining them in a latent state. Blocks TRAF2 binding to LMP1 and inhibits LMP1-mediated NF-kappa-B activation. Negatively regulates NF-kappaB signaling and cell survival upon DNA damage. Plays a role as an adapter to assemble ZC3H12A, USP10 in a deubiquitination complex which plays a negative feedback response to attenuate NF-kappaB activation through the deubiquitination of IKBKG or TRAF6 in response to interleukin-1-beta (IL1B) stimulation or upon DNA damage. Promotes UBP10-induced deubiquitination of TRAF6 in response to DNA damage. May control negatively TRAF2-mediated NF-kappa-B activation signaled by CD40, TNFR1 and TNFR2. Essential for the efficient induction of IRF-dependent transcription following infection with Sendai virus. {ECO:0000250 UniProtKB:Q92844}.</p>
Molecular Weight:	50.9 kDa
UniProt:	P70347
Pathways:	p53 Signaling , TLR Signaling , Activation of Innate immune Response

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p>

Application Details

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months