

Datasheet for ABIN3121679

Gasdermin C2 Protein (GSDMC2) (AA 1-480) (Strep Tag)



Overview

Quantity:	1 mg
Target:	Gasdermin C2 (GSDMC2)
Protein Characteristics:	AA 1-480
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This Gasdermin C2 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details	
Brand:	AliCE®
Sequence:	MGYSFDRASK DVVKKLQGRD LRPVECLSDA TKFRLFHILQ ETPRSGWETE DIPVGFTLLD
	LLEPNFPVPE PEVSAPKPFI HVQSTDLEAN LNVADIARGG VGYVGYGGYN IEVQSTSIPN
	PKLEILQNRK LLDNLPTFMK FCRMERKNLY VVTEAYEVSK DTMLTGLSSV NLSVKGFFKQ
	LFKVRGKAGR SEKYSIPIPK GSVLAYKKQQ LVIENNTCVI LPSATKKKMT FPGTPKYASA
	SEPTEIYRTE LQGLWINDIV PIGRIQEPAH LDFMCLQNEV YKQTEQLAEL SKGVQEVVLS
	SILSMLYEGD RKVLYDLMNM LELNQLGHMD GPGGKILDEL RKDSSNPCVD LKDLILYLLQ
	ALMVLSDSQL NLLAQSVEMG ILPHQVELVK SILQPNFKYP WNIPFTLQPQ LLAPLQGEGL
	AITYELLEEC GLKMELNNPR STWDLEAKMP LSALYGSLSF LQQLRKANSS SKPSLRPGYI
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	Gasdermin C2 (GSDMC2)
Alternative Name:	Gsdmc2
Background:	Gasdermin-C2 [Cleaved into: Gasdermin-C2, N-terminal (GSDMC2-NT), Gasdermin-C2, C-terminal (GSDMC2-CT)],FUNCTION: [Gasdermin-C2]: This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-C2, N-terminal) binds to membranes and forms pores, triggering pyroptosis. {ECO:0000269 PubMed:34290141}., FUNCTION: [Gasdermin-C2, N-terminal]: Pore-forming protein that causes membrane permeabilization and pyroptosis in response to type-2 immunity (PubMed:34290141). Produced by the cleavage of gasdermin-D in response to type-2 immunity following worm infection (PubMed:34290141). After cleavage, moves to the plasma membrane where it strongly binds to membrane inner leaflet lipids (By similarity). Homooligomerizes within the membrane and forms pores of 10-15 nanometers (nm) of inner diameter, triggering pyroptosis and lytic cell death in enterocytes (PubMed:34290141). {ECO:0000250 UniProtKB:Q9BYG8, ECO:0000269 PubMed:34290141}.
Molecular Weight:	54.0 kDa
UniProt: Application Details	Q2KHK6
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Application Notes: Comment:	as well. As the protein has not been tested for functional studies yet we cannot offer a

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months