

Datasheet for ABIN3121732

Gasdermin A2 Protein (GSDMA2) (AA 1-443) (Strep Tag)[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	Gasdermin A2 (GSDMA2)
Protein Characteristics:	AA 1-443
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This Gasdermin A2 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Brand:	AliCE®
Sequence:	MSMFEDVTRA LARQLNPRGD LTPLDSLIDF KRFHPFCLVL RKRKSTLFWG ARYVRTDYTL LDVLEPGSSP SDPTLLGNFS FKNMLDVRVE GDVEVPTMMK VKGTVGLSQS STLEVQMLSV APTALLENLHM ERKLSADHPF LKEMREYKQN LYVVMDEVKA KQEVTLKRAS NAISKFSLNL PSLGLQGSVN HKEAVTIPKG CVLAYRVRQL IYVGKDEWGI PYICTDNMPT FNPLCVLQRQ GSTVQMISGE MHEDFKTLKK EVQQETQVE KLSPVGRSSL LTSLSHLLGK KKELDLEQM LEGALDKGHE VTLEALPKDV LLLKDAMDAI LYFLGALTEL SEEQLKILVK SLENKVLVPVQ LKLVESILEQ NFLQDKEDVF PLRPDLLSSL GEEDQILTEA LVGLSGLEVQ RSGPQYTWNP DTCHNLALY AGLSLLHLLS RDS

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Product Details

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target: Gasdermin A2 (GSDMA2)

Alternative Name: Gsdma2

Background: Gasdermin-A2 (Gasdermin-2) [Cleaved into: Gasdermin-A2, N-terminal (GSDMA2-NT), Gasdermin-A2, C-terminal (GSDMA2-CT)],FUNCTION: [Gasdermin-A2]: This form constitutes the precursor of the pore-forming protein and acts as a sensor of infection: upon bacterial infection, specifically cleaved by some bacterial effector protein, releasing the N-terminal moiety (Gasdermin-A2, N-terminal) that binds to membranes and forms pores, triggering pyroptosis. {ECO:0000250|UniProtKB:Q9EST1}., FUNCTION: [Gasdermin-A2, N-terminal]: Pore-forming protein that causes membrane permeabilization and pyroptosis. Released upon cleavage of Gasdermin-A2, and binds to membrane inner leaflet lipids. Homooligomerizes within the membrane and forms pores of 10-15 nanometers (nm) of inner diameter, triggering pyroptosis. Binds to membrane inner leaflet lipids, such as phosphatidylinositol (4,5)-bisphosphate. {ECO:0000250|UniProtKB:Q96QA5}.

Molecular Weight: 49.8 kDa

UniProt: [Q32M21](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months