

Datasheet for ABIN3121748

## NHEJ1 Protein (AA 1-295) (Strep Tag)



[Go to Product page](#)

### Overview

Quantity:	250 µg
Target:	NHEJ1
Protein Characteristics:	AA 1-295
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NHEJ1 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB)

### Product Details

Brand:	AliCE®
Sequence:	<p>MEELEQDLLL QPRAWLQLAE NSLLAKVSIT KHGYALLISD LQQVWHEQVD TSVVSQRAKE  LNKRLTAPPA ALLCHLDEAL RPLFKDSAHP SKATFSCDRG EEGLILRVQS ELISGLPFSWH  FHCIPASSSL VSQHLLHPLM GVSLALQSHV RELAALLRMK DLEIQAYQES GAVLSRSRLK  TEPFEENSFL EQFMAEKLPE ACAVGDGKPF AMSLQSLYVA VTKQQIQARQ AHKDSGETQA  SSSTSPRGTD NQPEEPVSLs STLSEPEYEP VAASGPMHRA RLVKSKRKKP RGLFS</p> <p><b>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> </ul>
------------------	---

## Product Details

---

- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

## Target Details

---

Target:	NHEJ1
Alternative Name:	Nhej1 ( <a href="#">NHEJ1 Products</a> )

## Target Details

Background:	<p>Non-homologous end-joining factor 1 (Protein cernunnos) (XRCC4-like factor),FUNCTION: DNA repair protein involved in DNA non-homologous end joining (NHEJ), required for double-strand break (DSB) repair and V(D)J recombination (PubMed:17360556, PubMed:27601299, PubMed:27798842, PubMed:27601633, PubMed:27830975, PubMed:28051062, PubMed:29077092, PubMed:30017584). Plays a key role in NHEJ by promoting the ligation of various mismatched and non-cohesive ends (PubMed:17360556). Interacts with POLL (DNA polymerase lambda), promoting POLL recruitment to double-strand breaks (DSBs) and stimulation of the end-filling activity of POLL (By similarity). May act in concert with XRCC5-XRCC6 (Ku) to stimulate XRCC4-mediated joining of blunt ends and several types of mismatched ends that are non-complementary or partially complementary (PubMed:17360556). In some studies, has been shown to associate with XRCC4 to form alternating helical filaments that bridge DNA and act like a bandage, holding together the broken DNA until it is repaired (By similarity). Alternatively, it has also been shown that rather than forming filaments, a single NHEJ1 dimer interacts through both head domains with XRCC4 to promote the close alignment of DNA ends (By similarity). The XRCC4-NHEJ1/XLF subcomplex binds to the DNA fragments of a DSB in a highly diffusive manner and robustly bridges two independent DNA molecules, holding the broken DNA fragments in close proximity to one other (By similarity). The mobility of the bridges ensures that the ends remain accessible for further processing by other repair factors (By similarity). Binds DNA in a length-dependent manner (By similarity). {ECO:0000250 UniProtKB:A0A1L8ENT6, ECO:0000250 UniProtKB:Q9H9Q4, ECO:0000269 PubMed:17360556, ECO:0000269 PubMed:27601299, ECO:0000269 PubMed:27601633, ECO:0000269 PubMed:27798842, ECO:0000269 PubMed:27830975, ECO:0000269 PubMed:28051062, ECO:0000269 PubMed:29077092, ECO:0000269 PubMed:30017584}.</p>
-------------	---

Molecular Weight:	32.7 kDa
-------------------	----------

UniProt:	<a href="#">Q3KNJ2</a>
----------	------------------------

Pathways:	<a href="#">DNA Damage Repair</a>
-----------	-----------------------------------

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
--------------------	--

Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
----------	--

Application Details

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months