antibodies .- online.com





SGK196 Protein (AA 1-349) (rho-1D4 tag)



Image



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	1 mg
Target:	SGK196
Protein Characteristics:	AA 1-349
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SGK196 protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence:

MGQQHGTRNG LTHRELPRGV GLLLAMALMN VALYLCLDQL FISPGRSTAD SRRCPPGYFR
MGRMRNCSRW LSCEELRTEV RQLKRVGEGA VKRVFLSEWK EHKVALSRLT RLEMKEDFLH
GLQMLKSLQS EHVVTLVGYC EEDGTILTEY HPLGSLSNLE ETLNLSKYQD VNTWQHRLQL
AMEYVSIINY LHHSPLGTRV MCDSNDLPKT LSQYLLTSNF SIVANDLDAL PLVDHDSGVL
IKCGHRELHG DFVAPEQLWP YGEDTPFQDD LMPSYNEKVD IWKIPDVSSF LLGHVEGSDM
VRFHLFDIHK ACKSQIPAER PTAQNVLDAY QRVFHSLRDT VMSQTKEML

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Pomk Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Protein is endotoxin-free.
Grade:	Crystallography grade

Target Details

Target: SGK196

Target Details

Protein O-mannose kinase that specifically mediates phosphorylation at the 6-position of an O-mannose of the trisaccharide (N-acetylgalactosamine (GalNAc)-beta-1,3-N-acetylglucosamine (GlcNAc)-beta-1,4-mannose) to generate phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-1,3-N-acetylglucosamine-beta-1,4-(phosphate-6-)mannose). Phosphorylated O-mannosyl trisaccharide is a carbohydrate structure present in alphadystroglycan (DAG1), which is required for binding laminin G-like domain-containing extracellular proteins with high affinity. Only shows kinase activity when the GalNAc-beta-3-	
(GlcNAc)-beta-1,4-mannose) to generate phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-1,3-N-acetylglucosamine-beta-1,4-(phosphate-6-)mannose). Phosphorylated O-mannosyl trisaccharide is a carbohydrate structure present in alphadystroglycan (DAG1), which is required for binding laminin G-like domain-containing	
acetylgalactosamine-beta-1,3-N-acetylglucosamine-beta-1,4-(phosphate-6-)mannose). Phosphorylated O-mannosyl trisaccharide is a carbohydrate structure present in alphadystroglycan (DAG1), which is required for binding laminin G-like domain-containing	
Phosphorylated O-mannosyl trisaccharide is a carbohydrate structure present in alphadystroglycan (DAG1), which is required for binding laminin G-like domain-containing	
dystroglycan (DAG1), which is required for binding laminin G-like domain-containing	
extracellular proteins with high affinity. Only shows kinase activity when the GalNAc-beta-2-	
Characterial proteins with high affility. Only shows kinds activity when the dally actuals	
GlcNAc-beta-terminus is linked to the 4-position of O-mannose, suggesting that this	
disaccharide serves as the substrate recognition motif (By similarity). {ECO:0000250}.	
41.1 kDa Including tag.	
Q3TUA9	
In addition to the applications listed above we expect the protein to work for functional studies	
as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.	
Protein has not been tested for activity yet. In cases in which it is highly likely that the	
recombinant protein with the default tag will be insoluble our protein lab may suggest a higher	
molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible	
options with you in detail to assure that you receive your protein of interest.	
For Research Use only	
Liquid	
100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Avoid repeated freeze-thaw cycles.	
-80 °C	
Store at -80°C.	
Unlimited (if stored properly)	



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process