

Datasheet for ABIN3122620

MYST2 Protein (AA 1-613) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	MYST2
Protein Characteristics:	AA 1-613
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This MYST2 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Brand:	ALiCE®
Sequence:	<p>MAIGVVKRNA GSSSDGTEDS DFSTDLEHTD SSES DGTSRR SARVTRSSAR LSQSSQDSSP</p> <p>VRNLPSFGTE EPAYSTRRTV RSQQQPTPVT PKKYPLRQTR SSGSETEQVV DFSDRET KNT</p> <p>ADHDESPRPT PTGNAPSSSES DIDISSPNVS HDESI AKDMS LKDSGSDLSH RPKRRRFHES</p> <p>YNFNMKCPTP GCNSLGHLTG KHERHFSISG CPLYHNLSAD ECKVRAQSRD KQIEERMLSH</p> <p>RQDDNNRHAT RHQAPTERQL RYKEKVAELR KKRNSGLSKE QKEK YMEHRQ TYGNTREPLL</p> <p>ENLTSEYDLD LFRRAQARAS EDLEKLRLQG QITEGSNMIK TIAFGRYELD TWYHSPYPEE</p> <p>YARLGRLYMC EFCLKYMKSQ TILRRHMAK VVKHPPGDEI YRKG SISVFE VDGKKNKIYC</p> <p>QNLCLLAKLF LDHKTLYYDV EPFLFYVMTE ADNTGCHLIG YFSKEKNSFL NYNVSCILTM</p> <p>PQYMRQGYGK MLIDFSYLLS KVEEKVGSPE RPLSDLGLIS YRSYWK EVLL RYLHNFQGKE</p> <p>ISIKEISQET AVNPVDIVST LQALQMLKYW KGKHLVLKRQ DLIDEWIAKE AKRSNSNKT M</p> <p>DPSCCLKWTPP KGT</p>

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Product Details

Grade: custom-made

Target Details

Target: MYST2

Alternative Name: Kat7 ([MYST2 Products](#))

Background: Histone acetyltransferase KAT7 (EC 2.3.1.48) (Histone acetyltransferase binding to ORC1) (Lysine acetyltransferase 7) (MOZ, YBF2/SAS3, SAS2 and TIP60 protein 2) (MYST-2),FUNCTION: Catalytic subunit of histone acetyltransferase HBO1 complexes, which specifically mediate acetylation of histone H3 at 'Lys-14' (H3K14ac), thereby regulating various processes, such as gene transcription, protein ubiquitination, immune regulation, stem cell pluripotent and self-renewal maintenance and embryonic development (PubMed:21753189, PubMed:21149574, PubMed:23319590, PubMed:27733580, PubMed:31827282). Some complexes also catalyze acetylation of histone H4 at 'Lys-5', 'Lys-8' and 'Lys-12' (H4K5ac, H4K8ac and H4K12ac, respectively), regulating DNA replication initiation, regulating DNA replication initiation (By similarity). Specificity of the HBO1 complexes is determined by the scaffold subunit: complexes containing BRPF scaffold (BRPF1, BRD1/BRPF2 or BRPF3) direct KAT7/HBO1 specificity towards H3K14ac, while complexes containing JADE (JADE1, JADE2 and JADE3) scaffold direct KAT7/HBO1 specificity towards histone H4 (By similarity). H3K14ac promotes transcriptional elongation by facilitating the processivity of RNA polymerase II (PubMed:31827282). Acts as a key regulator of hematopoiesis by forming a complex with BRD1/BRPF2, directing KAT7/HBO1 specificity towards H3K14ac and promoting erythroid differentiation (By similarity). H3K14ac is also required for T-cell development (PubMed:27733580). KAT7/HBO1-mediated acetylation facilitates two consecutive steps, licensing and activation, in DNA replication initiation: H3K14ac facilitates the activation of replication origins, and histone H4 acetylation (H4K5ac, H4K8ac and H4K12ac) facilitates chromatin loading of MCM complexes, promoting DNA replication licensing (By similarity). Acts as a positive regulator of centromeric CENPA assembly: recruited to centromeres and mediates histone acetylation, thereby preventing centromere inactivation mediated by SUV39H1, possibly by increasing histone turnover/exchange (By similarity). Involved in nucleotide excision repair: phosphorylation by ATR in response to ultraviolet irradiation promotes its localization to DNA damage sites, where it mediates histone acetylation to facilitate recruitment of XPC at the damaged DNA sites (By similarity). Acts as an inhibitor of NF-kappa-B independently of its histone acetyltransferase activity (By similarity). {ECO:0000250|UniProtKB:O95251, ECO:0000269|PubMed:21149574, ECO:0000269|PubMed:21753189, ECO:0000269|PubMed:23319590, ECO:0000269|PubMed:27733580,

Target Details

ECO:0000269|PubMed:31827282}. FUNCTION: Plays a central role in the maintenance of leukemia stem cells in acute myeloid leukemia (AML) (PubMed:31827282). Acts by mediating acetylation of histone H3 at 'Lys-14' (H3K14ac), thereby facilitating the processivity of RNA polymerase II to maintain the high expression of key genes, such as HOXA9 and HOXA10 that help to sustain the functional properties of leukemia stem cells (PubMed:31827282). {ECO:0000269|PubMed:31827282}.

Molecular Weight: 70.6 kDa

UniProt: [Q5SVQ0](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Handling

Expiry Date: 12 months