

Datasheet for ABIN3122748

NRF2 Protein (AA 1-597) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	NRF2 (NFE2L2)
Protein Characteristics:	AA 1-597
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NRF2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MMDLELPPPG LQSQQDMDLI DILWRQDIDL GVSREVFDFS QRQKDYELEK QKKLEKERQE QLQKEQEKAFAQFQLDEET GEFLPIQPAQ HIQTDTSGSA SYSQVAHIPK QDALYFEDCM QLLAETFPFV DDHESLALDI PSHAESSVFT APHQAQSLNS SLEAAMTDLS SIEQDMEQVW QELFSIPELQ CLNTENKQLA DTTAVPSPEA TLTEMDSNYH FYSSISSLEK EVGNCGPHFL HGFEDSFSSI LSTDDASQLT SLDSNPTLNT DFGDEFYSAF IAEPDGGGSM PSSAAISQSL SELLDGTIEG CDLSLCKAFN PKHAEGTMEF NDSDSGISLN TSPSRASPEH SVESSYGDGP PPGFSDSEME ELDSAPGSVK QNGPKAQPAH SPGDTVQPLS PAQGHSA PMR ESQCEN TTKK EVPVSPGHQK APFTKDKHSS RLEAHLTRDE LRAKALHIPF PVEKIINLPV DDFNEMMSKE QFNEAQLALI RDIRRGKNK VAAQNCRKRK LENIVELEQD LGHLKDEREK LLREKGENDR NLHLLKRRLS TLYLEVFSML RDEDGKPYSP SEYSLQQTRD GNVFLVPKSK KPDTKKK</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression</p>

system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	NRF2 (NFE2L2)
Alternative Name:	Nfe2l2 (NFE2L2 Products)
Background:	<p>Nuclear factor erythroid 2-related factor 2 (NF-E2-related factor 2) (NFE2-related factor 2) (Nuclear factor, erythroid derived 2, like 2),FUNCTION: Transcription factor that plays a key role in the response to oxidative stress: binds to antioxidant response (ARE) elements present in the promoter region of many cytoprotective genes, such as phase 2 detoxifying enzymes, and promotes their expression, thereby neutralizing reactive electrophiles (PubMed:9240432, PubMed:9887101, PubMed:12032331, PubMed:14517554, PubMed:31398338). In normal conditions, ubiquitinated and degraded in the cytoplasm by the BCR(KEAP1) complex (PubMed:15282312, PubMed:15367669, PubMed:15581590). In response to oxidative stress, electrophile metabolites inhibit activity of the BCR(KEAP1) complex, promoting nuclear accumulation of NFE2L2/NRF2, heterodimerization with one of the small Maf proteins and binding to ARE elements of cytoprotective target genes (PubMed:12032331). The NFE2L2/NRF2 pathway is also activated in response to selective autophagy: autophagy promotes interaction between KEAP1 and SQSTM1/p62 and subsequent inactivation of the BCR(KEAP1) complex, leading to NFE2L2/NRF2 nuclear accumulation and expression of cytoprotective genes (PubMed:20421418, PubMed:20173742). May also be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region (By similarity). Also plays an important role in the regulation of the innate immune response. It is a critical regulator of the innate immune response and survival during sepsis by maintaining redox homeostasis and restraint of the dysregulation of pro-inflammatory signaling pathways like MyD88-dependent and -independent and TNF-alpha signaling (PubMed:16585964). Suppresses macrophage inflammatory response by blocking pro-inflammatory cytokine transcription and the induction of IL6 (PubMed:27211851). Binds to the proximity of pro-inflammatory genes in macrophages and inhibits RNA Pol II recruitment. The inhibition is independent of the Nrf2-binding motif and reactive oxygen species level (PubMed:27211851). Represses antiviral cytosolic DNA sensing by suppressing the expression of the adapter protein STING1 and decreasing responsiveness to STING1 agonists while increasing susceptibility to infection with DNA viruses (By similarity).</p> <p>{ECO:0000250 UniProtKB:Q16236, ECO:0000269 PubMed:12032331, ECO:0000269 PubMed:14517554, ECO:0000269 PubMed:15282312, ECO:0000269 PubMed:15367669, ECO:0000269 PubMed:15581590, ECO:0000269 PubMed:16585964, ECO:0000269 PubMed:20173742, ECO:0000269 PubMed:20421418, ECO:0000269 PubMed:27211851, ECO:0000269 PubMed:31398338, ECO:0000269 PubMed:9240432,</p>

Target Details

	ECO:0000269 PubMed:9887101}.
Molecular Weight:	66.9 kDa
UniProt:	Q60795
Pathways:	ER-Nucleus Signaling , Negative Regulation of intrinsic apoptotic Signaling

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	<p>The buffer composition is at the discretion of the manufacturer.</p> <p>Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.</p>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months