.-online.com antibodies

Datasheet for ABIN3122819 GNGT1 Protein (AA 2-71) (His tag)



Overview	
Quantity:	1 mg
Target:	GNGT1
Protein Characteristics:	AA 2-71
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This GNGT1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)
Product Details	
Sequence:	PVINIEDLTE KDKLKMEVDQ LKKEVTLERM MVSKCCEEVR DYIEERSGED PLVKGIPEDK
	NPFKELKGGC
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Mouse Gngt1 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3122819 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

cannot be expressed or purified. In the unlikely event that the protein cannot be expressed or purified w (other companies might charge you for any performed steps in the exp custom-made proteins, e.g. fees might apply for the expression plasm experiments or purification optimization). When you order this made-to-order protein you will only pay upon rece folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cel different His-tag capture materials: high yield, EDTA resistant, or DT	oression process for id, the first expression vival of the correctly that our experienced otein you ordered. absorbance at 280nm.
(other companies might charge you for any performed steps in the exp custom-made proteins, e.g. fees might apply for the expression plasm experiments or purification optimization).When you order this made-to-order protein you will only pay upon rece folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro- The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer.Purification:Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	oression process for id, the first expression vival of the correctly that our experienced otein you ordered. absorbance at 280nm.
custom-made proteins, e.g. fees might apply for the expression plasm experiments or purification optimization).When you order this made-to-order protein you will only pay upon rece folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer.The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification:Purification:Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	id, the first expression ival of the correctly that our experienced otein you ordered. absorbance at 280nm.
experiments or purification optimization). When you order this made-to-order protein you will only pay upon rece folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro- The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cel	tival of the correctly that our experienced otein you ordered. absorbance at 280nm.
When you order this made-to-order protein you will only pay upon rece folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro- The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification:Purification:Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	that our experienced otein you ordered. absorbance at 280nm.
folded protein. With no financial risk on your end you can rest assured protein experts will do everything to make sure that you receive the pro- The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer.The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	that our experienced otein you ordered. absorbance at 280nm.
protein experts will do everything to make sure that you receive the product The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification:Purification:Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	otein you ordered. absorbance at 280nm.
The concentration of our recombinant proteins is measured using the The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	absorbance at 280nm.
The protein's absorbance will be measured in several dilutions and is r specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	
specific reference buffer. The concentration of the protein is calculated using its specific absorp the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cel	1
The concentration of the protein is calculated using its specific absorption the Expasy's protparam tool to determine the absorption coefficient of Purification: Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell	neasured against its
the Expasy's protparam tool to determine the absorption coefficient ofPurification:Two step purification of proteins expressed in bacterial culture:1. In a first purification step, the protein is purified from the cleared cell	
Purification:Two step purification of proteins expressed in bacterial culture:1. In a first purification step, the protein is purified from the cleared cell	otion coefficient. We use
1. In a first purification step, the protein is purified from the cleared cel	f each protein.
different His-tag capture materials: high yield, EDTA resistant, or DT	ll lysate using three
	T resistant. Eluate
fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to	second purification stop
through size exclusion chromatography. Eluate fractions are analyze Western blot.	
Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography an	nd Western Blot.
Sterility: 0.22 µm filtered	
Endotoxin Level: Endotoxin has not been removed. Please contact us if you require end	otoxin removal.
Grade: Crystallography grade	

Target Details

Target:	GNGT1
Alternative Name:	Gngt1 (GNGT1 Products)
Background:	Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction.
Molecular Weight:	9.1 kDa Including tag.
UniProt:	Q61012

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN3122819 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Pathways:	Myometrial Relaxation and Contraction, Regulation of G-Protein Coupled Receptor Protein Signaling, Phototransduction
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)