antibodies .-online.com

Datasheet for ABIN3123405 SRD5A1 Protein (AA 1-255) (Strep Tag)



Overview

| 1 mg |
|--|
| SRD5A1 |
| AA 1-255 |
| Mouse |
| Tobacco (Nicotiana tabacum) |
| Recombinant |
| This SRD5A1 protein is labelled with Strep Tag. |
| Western Blotting (WB), SDS-PAGE (SDS), ELISA |
| |
| MELDELCLLD ALVYLEGFLA FVAFVGLQMV GSSYGRYSSQ WSGRRVPARP AWFLQELPSM |
| AWPLYECIRP AAARLGNLPN RVLLAMFLIH YVQRTLVFPV LIRGGKPTLL FTFVLAFLFC |
| TLNGYLQSRY LSQFAVYAED WVTHPCFLTG FALWLVGMVI NIHSDHILRN LRKPGETGYK |
| IPRGGLFEYV SSANYFGELV EWCGFALASW SLQGVVFALF TLCALFTRAR QHHQWYLEKF |
| EDYPKTRKIL IPFLL |
| Sequence without tag. The proposed Strep-Tag is based on experience s with the expression |
| system, a different complexity of the protein could make another tag necessary. In case you |
| have a special request, please contact us. |
| Key Benefits: |
| Made in Germany - from design to production - by highly experienced protein experts. |
| • Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification. |
| |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3123405 | 05/07/2024 | Copyright antibodies-online. All rights reserved.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

| Purification:Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®): 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) | | |
|--|------------------|--|
| capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. Purity: ≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. | Purification: | |
| | | capture material. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and |
| Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) | Purity: | ≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
| | Endotoxin Level: | Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3123405 | 05/07/2024 | Copyright antibodies-online. All rights reserved.

| Target: | SRD5A1 |
|---------------------------------------|---|
| Alternative Name: | Srd5a1 (SRD5A1 Products) |
| Background: | 3-oxo-5-alpha-steroid 4-dehydrogenase 1 (EC 1.3.1.22) (SR type 1) (Steroid 5-alpha-reductase 1) (S5AR 1),FUNCTION: Converts testosterone into 5-alpha-dihydrotestosterone and progesterone or corticosterone into their corresponding 5-alpha-3-oxosteroids. It plays a centra role in sexual differentiation and androgen physiology. {ECO:0000250 UniProtKB:P24008}. |
| Molecular Weight: | 29.3 kDa |
| UniProt: | Q68FF9 |
| Pathways: | Metabolism of Steroid Hormones and Vitamin D, Steroid Hormone Biosynthesis, Response to Growth Hormone Stimulus, C21-Steroid Hormone Metabolic Process |
| Application Details | |
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. |
| Comment: | ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for |
| | protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein! |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us. |
| · · · · · · · · · · · · · · · · · · · | |

Handling Advice: Avoid repeated freeze-thaw cycles.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3123405 | 05/07/2024 | Copyright antibodies-online. All rights reserved.

Handling

| Storage: | -80 °C |
|------------------|--------------------------------|
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |