

Datasheet for ABIN3123586

CCDC111 Protein (AA 1-537) (Strep Tag)



Overview

Quantity:	250 μg
Target:	CCDC111
Protein Characteristics:	AA 1-537
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCDC111 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Dotaile

Product Details	
Brand:	AliCE®
Sequence:	MLRKWEARVK QIEERASHYE RKPLSSVYRP RLAKPEEPSS IWKLFHRQNQ AFNFVKSCKE
	SVHVFALECK RGNGQRIYLV TSYAQLWFYY KTRKTLLHCY EVIPENAVCK LYFDLEFNKL
	ANPGADGKMM VALLIQHVCK ALEEFYNVQC SAEDVFNLDS STEEKFSRHL IFQLHNVAFK
	DNRHAGNFVR KILQPALHLI AEDDEAKVPE AVGQDASGFS VTPLKQEISE AREKVGLPKQ
	CDPDLSFLVV KNHMGEKCLF VDLGVYTKNR NFRLYQSSKI GKCVSLEVAE DNRFIPKQSK
	DISEENQYFL SSLVSNVRFS DTLRVLTCHP SQTKRKRAEC FNSTGTSVES IEGFQGSPYP
	EVDQFVLSLV NKHDIKGGIR RWNYFFPEEL LVYDICKYRW CENIGRAHKS NNIMILVDLK
	NEVWYQKCHD PVCKAQNFKS TCSPLPTEVS LLFLLKDEDF TSGETDDTST SLTKDSQTPP
	SCNLSAGGLS AAAWDDEDDA LFLEATEDAE FADAADKSLG SMDDIPDELI IEALQNS
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	CCDC111
Alternative Name:	Primpol (CCDC111 Products)
Background:	DNA-directed primase/polymerase protein (EC 2.7.7),FUNCTION: DNA primase and DNA
	polymerase required to tolerate replication-stalling lesions by bypassing them
	(PubMed:26926109, PubMed:29073063). Required to facilitate mitochondrial and nuclear
	replication fork progression by initiating de novo DNA synthesis using dNTPs and acting as an
	error-prone DNA polymerase able to bypass certain DNA lesions (By similarity). Shows a high
	capacity to tolerate DNA damage lesions such as 80xoG and abasic sites in DNA (By similarity
	Provides different translesion synthesis alternatives when DNA replication is stalled: able to
	synthesize DNA primers downstream of lesions, such as ultraviolet (UV) lesions, R-loops and C
	quadruplexes, to allow DNA replication to continue (By similarity). Can also realign primers
	ahead of 'unreadable lesions' such as abasic sites and 6-4 photoproduct (6-4 pyrimidine-
	pyrimidinone), thereby skipping the lesion (By similarity). Also able to incorporate nucleotides
	opposite DNA lesions such as 80xoG, like a regular translesion synthesis DNA polymerase (By
	similarity). Also required for reinitiating stalled forks after UV damage during nuclear DNA
	replication (By similarity). Required for mitochondrial DNA (mtDNA) synthesis and replication,
	by reinitiating synthesis after UV damage or in the presence of chain-terminating nucleotides
	(PubMed:29073063). Prevents APOBEC family-mediated DNA mutagenesis by repriming
	downstream of abasic site to prohibit error-prone translesion synthesis (PubMed:26926109).
	Has non-overlapping function with POLH (By similarity). In addition to its role in DNA damage
	response, also required to maintain efficient nuclear and mitochondrial DNA replication in
	unperturbed cells (By similarity). {ECO:0000250 UniProtKB:Q96LW4,
	ECO:0000269 PubMed:26926109, ECO:0000269 PubMed:29073063}.
Molecular Weight:	61.3 kDa
UniProt:	Q6P1E7
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational

Application Details

modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months