

Datasheet for ABIN3123610

SMUG1 Protein (AA 1-279) (Strep Tag)



()	11/	\sim	r١.	ıi.	0	۱۸/	,
U	V	H	r٧	1	C	V۷	

Quantity:	1 mg
Target:	SMUG1
Protein Characteristics:	AA 1-279
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SMUG1 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details	
Brand:	AliCE®
Sequence:	MAASQTFPLG PTHEPASALM EPLPCTRSLA EGFLEEELRL NAELSQLQFP EPVGVIYNPV
	DYAWEPHRNY VTRYCQGPKE VLFLGMNPGP FGMAQTGVPF GEVNVVRDWL GVGGPVLTPP
	QEHPKRPVLG LECPQSEVSG ARFWGFFRTL CGQPQVFFRH CFVHNLCPLL FLAPSGRNLT
	PAELPAKQRE QLLSICDAAL CRQVQLLGVR LVVGVGRLAE QRARRALAGL TPEVQVEGLL
	HPSPRSAQAN KGWEAAARER LQELGLLPLL TDEGSARPT
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:
	Made in Germany - from design to production - by highly experienced protein experts.

Alternative Name:

- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

Smug1 (SMUG1 Products)

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	SMUG1

Target Details

Background:	Single-strand selective monofunctional uracil DNA glycosylase (EC 3.2.2),FUNCTION:
	Recognizes base lesions in the genome and initiates base excision DNA repair. Acts as a
	monofunctional DNA glycosylase specific for uracil (U) residues in DNA with a preference for
	single-stranded DNA substrates. The activity is greater toward mismatches (U/G) compared to
	matches (U/A). Excises uracil (U), 5-formyluracil (fU) and uracil derivatives bearing an oxidized
	group at C5 [5-hydroxyuracil (hoU) and 5-hydroxymethyluracil (hmU)] in ssDNA and dsDNA, but
	not analogous cytosine derivatives (5-hydroxycytosine and 5-formylcytosine), nor other
	oxidized bases. The activity is damage-specific and salt-dependent. The substrate preference is
	the following: ssDNA > dsDNA (G pair) = dsDNA (A pair) at low salt concentration, and dsDNA
	(G pair) > dsDNA (A pair) > ssDNA at high salt concentration.
	{ECO:0000250 UniProtKB:Q53HV7}.
Molecular Weight:	30.7 kDa
UniProt:	Q6P5C5
Pathways:	DNA Damage Repair
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Handling

Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months