

Datasheet for ABIN3125921 **FEM1C Protein (AA 1-617) (Strep Tag)**



Overview

Quantity:	250 μg
Target:	FEM1C
Protein Characteristics:	AA 1-617
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This FEM1C protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Brand:	AliCE®
Sequence:	MDLKTAVFNA ARDGKLRLLT KLLASKSKAE VSSLISEKTN GATPLLMAAR YGHLDMVEFL
	LEQCSASIEV GGSVNFDGET IEGAPPLWAA SAAGHLKVVQ SLLNHGASVN NTTLTNSTPL
	RAACFDGHLE IVKYLVEHKA DLEVSNRHGH TCLMISCYKG HKEIAQYLLE KGADVNRKSV
	KGNTALHDCA ESGSLDIMKM LLMYCAKMEK DGYGMTPLLS ASVTGHTNIV DFLTHHAQTS
	KTERINALEL LGATFVDKKR DLLGALKYWK KAMNMRYSDR TNIISKPVPQ TLIMAYDYAK
	EVNSAEELEG LIADPDEMRM QALLIRERIL GPSHPDTSYY IRYRGAVYAD SGNFKRCINL
	WKYALDMQQS NLDPLSPMTA SSLLSFAELF SFMLQDRAKG LLGTTVTFDD LMGILCKSVL
	EIERAIKQTQ CPADPLQLNK ALSIILHLIC LLEKVPCTVE QDHFKKQTIY RFLKLHPRGK
	NNFSPLHLAV DKNTTCVGRY PVCKFPSLQV TAILIECGAD VNVRDSDDNS PLHIAALNNH
	PDIMNLLIKS GAHFDATNLH KQTASDLLDE KEIAKNLIQP INHTTLQCLA ARVIVNHRIY
	YKGNIPEKLE TFVSLHR

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

custom-made

Fem1c (FEM1C Products)

Protein fem-1 homolog C (FEM1c) (FEM1-gamma), FUNCTION: Substrate-recognition component of a Cul2-RING (CRL2) E3 ubiquitin-protein ligase complex of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation. The C-degron recognized by the DesCEND pathway is usually a motif of less than ten residues and can be present in full-length proteins, truncated proteins or proteolytically cleaved forms. The CRL2(FEM1C) complex specifically recognizes proteins with an arginine at the C-terminus: recognizes and binds proteins ending with -Lys/Arg-Xaa-Arg and -Lys/Arg-Xaa-Xaa-Arg Cdegrons, such as SIL1 or OR51B2, leading to their ubiquitination and degradation. The CRL2(FEM1C) complex mediates ubiquitination and degradation of truncated MSRB1/SEPX1 selenoproteins produced by failed UGA/Sec decoding. {ECO:0000250|UniProtKB:Q96JP0}.

Molecular Weight: 68.6 kDa

UniProt: Q8CEF1

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months