

Datasheet for ABIN3126062

## SPHK1 Protein (AA 1-382) (Strep Tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	SPHK1
Protein Characteristics:	AA 1-382
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SPHK1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

### Product Details

Brand:	AliCE®
Sequence:	<p>MEPVECPRGL LPRPCRVLVL LNPQGGKGKA LQLFQSRVQP FLEEAEITFK LILTERKNHA  RELVCAEELG HWDALAVMSG DGLMHEVVNG LMERPDWETA IQKPLCSLPG GSGNALAASV  NHYAGYEQVT NEDLLINCTL LLCRRRLSPM NLLSLHTASG LRLYSVL SLS WGFVADV DLE  SEKYRRLGEI RFTVGTFFRL ASLRIYQGQL AYLPVGTVAS KRPASTLVQK GPVDTHLVPL  EEPVPSHWTV VPEQDFVLVL VLLHTHLSSE LFAAPMGRCE AGVMHLFYVR AGVSRAALLR  LFLAMQKGKH MELDCPYLVH VPVVAFRLEP RSQRGVFSVD GELMVCEAVQ GQVHPNYLWM  VCGSRDAPSG RDSRRGPPPE EP</p> <p><b>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>
Characteristics:	Key Benefits:

## Product Details

---

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

## Target Details

---

Target:	SPHK1
---------	-------

---

## Target Details

Alternative Name: Sphk1 ([SPHK1 Products](#))

Background: Sphingosine kinase 1 (SK 1) (SPK 1) (EC 2.7.1.91) (Acetyltransferase SPHK1) (EC 2.3.1.-),FUNCTION: Catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (SPP), a lipid mediator with both intra- and extracellular functions (PubMed:17346996, PubMed:21084291, PubMed:25417698, PubMed:29662056, PubMed:33334894). Also acts on D-erythro-sphingosine and to a lesser extent sphinganine, but not other lipids, such as D,L-threo-dihydrosphingosine, N,N-dimethylsphingosine, diacylglycerol, ceramide, or phosphatidylinositol (PubMed:9726979). In contrast to proapoptotic SPHK2, has a negative effect on intracellular ceramide levels, enhances cell growth and inhibits apoptosis (PubMed:16118219). Involved in the regulation of inflammatory response and neuroinflammation. Via the product sphingosine 1-phosphate, stimulates TRAF2 E3 ubiquitin ligase activity, and promotes activation of NF-kappa-B in response to TNF signaling (By similarity). In response to TNF and in parallel to NF-kappa-B activation, negatively regulates RANTES induction through p38 MAPK signaling pathway (By similarity). Involved in endocytic membrane trafficking induced by sphingosine, recruited to dilate endosomes, also plays a role on later stages of endosomal maturation and membrane fusion independently of its kinase activity (PubMed:27806293, PubMed:28049734). In Purkinje cells, seems to be also involved in the regulation of autophagosome-lysosome fusion upon VEGFA (PubMed:25417698). {ECO:0000250|UniProtKB:Q9NYA1, ECO:0000269|PubMed:16118219, ECO:0000269|PubMed:21084291, ECO:0000269|PubMed:25417698, ECO:0000269|PubMed:27806293, ECO:0000269|PubMed:28049734, ECO:0000269|PubMed:33334894, ECO:0000269|PubMed:9726979, ECO:0000305|PubMed:29662056}., FUNCTION: Has serine acetyltransferase activity on PTGS2/COX2 in an acetyl-CoA dependent manner. The acetyltransferase activity increases in presence of the kinase substrate, sphingosine. During neuroinflammation, through PTGS2 acetylation, promotes neuronal secretion of specialized preresolving mediators (SPMs), especially 15-R-lipoxin A4, which results in an increase of phagocytic microglia. {ECO:0000269|PubMed:29662056}.

Molecular Weight: 42.4 kDa

UniProt: [Q8CI15](#)

Pathways: [VEGF Signaling](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies

## Application Details

---

as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

### Comment:

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

---

### Restrictions:

For Research Use only

## Handling

---

### Format:

Liquid

---

### Buffer:

The buffer composition is at the discretion of the manufacturer.

Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

---

### Handling Advice:

Avoid repeated freeze-thaw cycles.

---

### Storage:

-80 °C

---

### Storage Comment:

Store at -80°C.

---

### Expiry Date:

12 months