-online.com antibodies

## Datasheet for ABIN3126175 CDC42EP2 Protein (AA 2-214) (His tag)



Overview

Overview	
Quantity:	1 mg
Target:	CDC42EP2
Protein Characteristics:	AA 2-214
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CDC42EP2 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)
Product Details	
Sequence:	STKVPIYLKR GSRKGKKEKL RDLLSSDMIS PPLGDFRHTI HIGSGGGDDM FGDISFLQGK
	FHLLPGTAVE EAEEDGSFDL PFQFTRTTTV CGRELPDGLS PLLKNAISLP VIGGPQALTL
	PTAQAPPKPP RLHLESPQPS PQPSPQGAGN VDVWRIPEAG SPHNGMSPEP EAEEPFLSHA
	SSLLSLHVDL GPSILDDVLQ IMDHDLGRVQ IPT
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	<ul> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> <li>Mouse Cdc42ep2 Protein (raised in Insect Cells) purified by multi-step, protein-specific</li> </ul>
	process to ensure crystallization grade.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3126175 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.In the unlikely event that the protein cannot be expressed or purified we do not charge anyth (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.		
cannot be expressed or purified.In the unlikely event that the protein cannot be expressed or purified we do not charge anyth (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We u the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE 3. Purity:Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility.0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		The big advantage of ordering our made-to-order proteins in comparison to ordering custom
In the unlikely event that the protein cannot be expressed or purified we do not charge anyth (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We u the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE actions are analyzed by SDS-PAGE.Purify:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Protein is endotoxin free.		made proteins from other companies is that there is no financial obligation in case the protein
(other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We u the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		cannot be expressed or purified.
custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization). When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We ut the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification set analyzed by SDS-PAGE. 3. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Protein is endotoxin free.		In the unlikely event that the protein cannot be expressed or purified we do not charge anything
experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We ut the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		(other companies might charge you for any performed steps in the expression process for
When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression
folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We ut the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Protein is endotoxin free.		experiments or purification optimization).
<ul> <li>protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.</li> <li>Purification: Two step purification of proteins expressed in baculovirus infected SF9 insect cells:         <ol> <li>In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> </ol> </li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> </ul>		When you order this made-to-order protein you will only pay upon receival of the correctly
The concentration of our recombinant proteins is measured using the absorbance at 280nm         The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.         The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells:         1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.         2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.         Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.		folded protein. With no financial risk on your end you can rest assured that our experienced
The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		protein experts will do everything to make sure that you receive the protein you ordered.
specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		The concentration of our recombinant proteins is measured using the absorbance at 280nm.
The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		The protein's absorbance will be measured in several dilutions and is measured against its
the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.		specific reference buffer.
Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells:1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 μm filteredEndotoxin Level:Protein is endotoxin free.		The concentration of the protein is calculated using its specific absorption coefficient. We use
<ul> <li>In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> </ul>		the Expasy's protparam tool to determine the absorption coefficient of each protein.
<ul> <li>different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> </ul>	Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
<ul> <li>2. Protein containing fractions of the best purification are subjected to second purification s through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> </ul>		different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate
Sterility:     0.22 µm filtered       Endotoxin Level:     Protein is endotoxin free.		2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and
Endotoxin Level: Protein is endotoxin free.	Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
	Sterility:	0.22 μm filtered
Grade: Crystallography grade	Endotoxin Level:	Protein is endotoxin free.
	Grade:	Crystallography grade
Target Details	Target Details	
Target: CDC42EP2	Target:	CDC42EP2
Alternative Name: Cdc42ep2 (CDC42EP2 Products)	Alternative Name:	Cdc42ep2 (CDC42EP2 Products)
Background: Probably involved in the organization of the actin cytoskeleton. May act downstream of CDC	Background:	Probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42

to induce actin filament assembly leading to cell shape changes. Induces pseudopodia

formation in fibroblasts in a CDC42-dependent manner (By similarity). {ECO:0000250}.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN3126175 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Molecular Weight:	23.8 kDa Including tag.
UniProt:	Q8JZX9
Pathways:	Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN3126175 | 04/30/2024 | Copyright antibodies-online. All rights reserved.

Unlimited (if stored properly)

Expiry Date: