

Datasheet for ABIN3126326 PTPLAD1 Protein (AA 1-149) (His tag)



Overview Quantity: 1 mg Target: PTPLAD1 Protein Characteristics: AA 1-149 Origin: Mouse Source: Escherichia coli (E. coli) Protein Type: Recombinant Purification tag / Conjugate: This PTPLAD1 protein is labelled with His tag. Application: ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys) Product Details Sequence: METQVLTPHV YWAQRHRELY LRVELSDVQN PAISITDNVL HFKAQGHGAK GDNVYEFHLE FLDLVKPEPA YRLTQRQVNI TVQKKGSHWW ERLTKQEKRP LFLAPDFDRW LDESDAEMEL RAKEEERLNK LRLEREGSPE TLTNLKKGY Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us. Characteristics: • Made in Germany - from design to production - by highly experienced protein experts. Mouse Hacd3 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our made-to-order proteins in comparison to ordering custom

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3126326 | 08/02/2024 | Copyright antibodies-online. All rights reserved.

	chain of long- and very long-chain fatty acids/VLCFAs per cycle. This enzyme catalyzes the dehydration of the 3-hydroxyacyl-CoA intermediate into trans-2,3-enoyl-CoA, within each cycle
Background:	Catalyzes the third of the four reactions of the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process, allows the addition of two carbons to the
Alternative Name:	Hacd3 (PTPLAD1 Products)
Target:	PTPLAD1
Target Details	
Grade:	Crystallography grade
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Sterility:	0.22 µm filtered
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
	 Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
	 In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
Purification:	Two step purification of proteins expressed in bacterial culture:
	the Expasy's protparam tool to determine the absorption coefficient of each protein.
	The concentration of the protein is calculated using its specific absorption coefficient. We use
	specific reference buffer.
	The protein's absorbance will be measured in several dilutions and is measured against its
	protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm.
	folded protein. With no financial risk on your end you can rest assured that our experienced
	When you order this made-to-order protein you will only pay upon receival of the correctly
	experiments or purification optimization).
	custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression
	(other companies might charge you for any performed steps in the expression process for
	In the unlikely event that the protein cannot be expressed or purified we do not charge anything
	cannot be expressed or purified.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN3126326 | 08/02/2024 | Copyright antibodies-online. All rights reserved.

	of fatty acid elongation. Thereby, it participates to the production of VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators. Involved in Rac1-signaling pathways leading to the modulation of gene expression. Promotes insulin receptor/INSR autophosphorylation and is involved in INSR internalization (By similarity). {ECO:0000250 UniProtKB:Q9P035}.
Molecular Weight:	18.7 kDa Including tag.
UniProt:	Q8K2C9
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C

Expiry Date:

Storage Comment:

Unlimited (if stored properly)

Store at -80°C.