
antibodies .- online.com





MGAT5 Protein (AA 1-740) (Strep Tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	MGAT5
Protein Characteristics:	AA 1-740
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This MGAT5 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MAFFSPWKLS SQKLGFFLVT FGFIWGMMLL HFTIQQRTQP ESSSMLREQI LDLSKRYIKA
LAEENRDVVD GPYAGVMTAY DLKKTLAVLL DNILQRIGKL ESKVDNLVNG TGANSTNSTT
AVPSLVSLEK INVADIINGV QEKCVLPPMD GYPHCEGKIK WMKDMWRSDP CYADYGVDGT
SCSFFIYLSE VENWCPRLPW RAKNPYEEAD HNSLAEIRTD FNILYGMMKK HEEFRWMRLR
IRRMADAWIQ AIKSLAEKQN LEKRKRKKIL VHLGLLTKES GFKIAETAFS GGPLGELVQW
SDLITSLYLL GHDIRISASL AELKEIMKKV VGNRSGCPTV GDRIVELIYI DIVGLAQFKK
TLGPSWVHYQ CMLRVLDSFG TEPEFNHASY AQSKGHKTPW GKWNLNPQQF YTMFPHTPDN
SFLGFVVEQH LNSSDIHHIN EIKRQNQSLV YGKVDSFWKN KKIYLDIIHT YMEVHATVYG
SSTKNIPSYV KNHGILSGRD LQFLLRETKL FVGLGFPYEG PAPLEAIANG CAFLNPKFNP
PKSSKNTDFF IGKPTLRELT SQHPYAEVFI GRPHVWTVDL NNREEVEDAV KAILNQKIEP
YMPYEFTCEG MLQRINAFIE KQDFCHGQVM WPPLSALQVK LAEPGQSCKQ VCQESQLICE
PSFFQHLNKE KDLLKYKVTC QSSELYKDIL VPSFYPKSKH CVFQGDLLLF SCAGAHPTHQ

RICPCRDFIK GQVALCKDCL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- · The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System

(ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Target Details

Target: MGAT5

Alternative Name: Mgat5 (MGAT5 Products)

Background:

Alpha-1,6-mannosylglycoprotein 6-beta-N-acetylglucosaminyltransferase A (EC 2.4.1.155) (Alpha-mannoside beta-1,6-N-acetylglucosaminyltransferase) (GlcNAc-T V) (GNT-V) (Mannoside acetylglucosaminyltransferase 5) (N-acetylglucosaminyl-transferase V) [Cleaved into: Secreted alpha-1,6-mannosylglycoprotein 6-beta-N-acetylglucosaminyltransferase A (Secreted beta-1,6-N-acetylglucosaminyltransferase V) (Secreted GNT-V)],FUNCTION: Catalyzes the addition of N-acetylglucosamine (GlcNAc) in beta 1-6 linkage to the alpha-linked mannose of biantennary N-linked oligosaccharides (PubMed:10700233, PubMed:14561752, PubMed:22715095). Catalyzes an important step in the biosynthesis of branched, complex-type N-glycans, such as those found on EGFR, TGFR (TGF-beta receptor) and CDH2 (PubMed:12122020, PubMed:10700233, PubMed:14561752, PubMed:15459394, PubMed:22715095). Via its role in the biosynthesis of complex N-glycans, plays an important role in the activation of cellular signaling pathways, reorganization of the actin cytoskeleton, cell-cell adhesion and cell migration (PubMed:10700233, PubMed:14561752, PubMed:15459394). MGAT5-dependent EGFR N-glycosylation enhances the interaction between EGFR and LGALS3 and thereby prevents rapid EGFR endocytosis and prolongs EGFR signaling (PubMed:15459394). Required for efficient interaction between TGFB1 and its receptor (PubMed:15459394). Enhances activation of intracellular signaling pathways by several types of growth factors, including FGF2, PDGF, IGF, TGFB1 and EGF (PubMed:15459394). MGAT5-dependent CDH2 N-glycosylation inhibits CDH2-mediated homotypic cell-cell adhesion and contributes to the regulation of downstream signaling pathways (PubMed:14561752). Promotes cell migration (PubMed:14561752, PubMed:15459394). Contributes to the regulation of the inflammatory response

(PubMed:11217864, PubMed:15459394). MGAT5-dependent TCR N-glycosylation enhances the interaction between TCR and LGALS3, limits agonist-induced TCR clustering, and thereby dampens TCR-mediated responses to antigens (PubMed:11217864). Required for normal leukocyte evasation and accumulation at sites of inflammation (PubMed:15459394). Inhibits attachment of monocytes to the vascular endothelium and subsequent monocyte diapedesis (By similarity). {ECO:0000250|UniProtKB:Q09328, ECO:0000269|PubMed:10700233, ECO:0000269|PubMed:11217864, ECO:0000269|PubMed:12122020,

ECO:0000269|PubMed:14561752, ECO:0000269|PubMed:15459394,

ECO:0000269|PubMed:22715095}., FUNCTION: [Secreted alpha-1,6-mannosylglycoprotein 6-beta-N-acetylglucosaminyltransferase A]: Promotes proliferation of umbilical vein endothelial cells and angiogenesis, at least in part by promoting the release of the growth factor FGF2 from the extracellular matrix. {ECO:0000250|UniProtKB:Q09328}.

Molecular Weight:

84.6 kDa

UniProt:

Q8R4G6

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:

Liquid

Handling

Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)