antibodies .- online.com





SETD3 Protein (AA 1-594) (Strep Tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	SETD3
Protein Characteristics:	AA 1-594
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SETD3 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MGKKSRVKTQ KSGTGATATV SPKEILNLTS ELLQKCSSPA PSPGKEWEEY TQIRALVEKI RKKQKGLSVT FDGKREDYFP DLMKWASENG ASVEGFEMVN FKEEGFGLRA TRDIKAEELF LWVPRKLLMT VESAKNSVLG PLYSQDRILQ AMGNIALAFH LLCERASPNS FWQPYIQTLP SEYDTPLYFE EEEVRCLQST QAIHDVFSQY KNTARQYAYF YKVIQTHPHA NKLPLKESFT YEDYRWAVSS VMTRQNQIPT EDGSRVTLAL IPLWDMCNHT NGLITTGYNL EDDRCECVAL QDFQAGDQIY IFYGTRSNAE FVIHSGFFFD NNSHDRVKIK LGVSKSDRLY AMKAEVLARA GIPTSSVFAL HSTEPPISAQ LLAFLRVFCM TEEELKEHLL GDSAIDRIFT LGNAEFPVSW DNEVKLWTFL EDRASLLLKT YKTTIEEDKI VLKNPDLSVR ATMAIKLRLG EKEILEKAVK SAAVNREYYR KHMEERAPLP RYEESDLGLL EGGVGDSRLP LVLRKLEEEA GVQESLSLTE TVSKVKAAEN GLVNGENLIP NGTRSENESL SPEESENVTG EESSGSMAKV KERL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.

	Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Target Details	
Target:	SETD3
Alternative Name:	Setd3 (SETD3 Products)
Background:	Actin-histidine N-methyltransferase (EC 2.1.1.85) (Endothelial differentiation inhibitory protein D10) (Protein-L-histidine N-tele-methyltransferase) (SET domain-containing protein 3),FUNCTION: Protein-histidine N-methyltransferase that specifically mediates 3-methylhistidine (tele-methylhistidine) methylation of actin at 'His-73' (PubMed:30626964). Histidine methylation of actin is required for smooth muscle contraction of the laboring uterus during delivery (PubMed:30626964). Does not have protein-lysine N-methyltransferase activity and probably only catalyzes histidine methylation of actin (PubMed:30626964). {ECO:0000269 PubMed:30626964}.
Molecular Weight:	67.2 kDa
UniProt:	Q91WC0
Pathways:	Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional

Application Details

	components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)