Datasheet for ABIN3127800 RNF144A Protein (AA 1-292) (Strep Tag)

antibodies.com



\sim						
()	۱ /	0	r\	/1	\sim	۱ ۸ <i>I</i>
()	v	-	I \	/ I	-	W
\sim		\sim	• •		\sim	•••

Overview	
Quantity:	1 mg
Target:	RNF144A
Protein Characteristics:	AA 1-292
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This RNF144A protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)
Product Details	
Sequence:	MTTARYRPTW DLALDPLVSC KLCLGEYPAE QMTTIAQCQC IFCTLCLKQY VELLIKEGLE
	TAISCPDAAC PKQGHLQENE IECMVAAEIM QRYKKLQFER EVLFDPCRTW CPASTCQAVC
	QLQDIGLQTP QLVQCKACDM EFCSACKARW HPGQGCPETM PITFLPGETS SAFKMEEGDA
	PIKRCPKCRV YIERDEGCAQ MMCKNCKHAF CWYCLESLDD DFLLIHYDKG PCRNKLGHSR
	ASVIWHRTQV VGIFAGFGLL LLVASPFLLL ATPFVLCCKC KCSKGDDDPL PT
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expressior
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:
	Made in Germany - from design to production - by highly experienced protein experts.
	Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3127800 | 05/08/2024 | Copyright antibodies-online. All rights reserved.

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein. The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®): 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)		
capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. Purity: ≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.	Purification:	
		capture material. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and
Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)	Purity:	\ge 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
	Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3127800 | 05/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	RNF144A
Alternative Name:	Rnf144a (RNF144A Products)
Background:	E3 ubiquitin-protein ligase RNF144A (EC 2.3.2.31) (RING finger protein 144A) (UbcM4- interacting protein 4) (Ubiquitin-conjugating enzyme 7-interacting protein 4),FUNCTION: E3 ubiquitin-protein ligase which accepts ubiquitin from E2 ubiquitin-conjugating enzymes UBE2L3 and UBE2L6 in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Mediates the ubiquitination and degradation of the DNA damage kinase PRKDC during DNA damage. Positively regulates DNA virus or exogenous cytosolic DNA-triggered innate immune response by mediating STING1 ubiquitination and increasing its 'Lys-6'-linked ubiquitination and translocation from the endoplasmic reticulum to the Golgi leading to downstream signaling pathways. Plays a positive role in EGF-dependent cell proliferation by prolonging EGF/EGFR signaling during EGF stimulation through EGFR ubiquitination. Increases ERK activity independently of EGFR signaling by promoting polyubiquitination and subsequent
Molecular Weight:	degradation of VRK3 in the cytosol. {ECO:0000250 UniProtKB:P50876}. 32.8 kDa
UniProt:	Q925F3

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.
	During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the
	mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce
	something that functions like a cell, but without the constraints of a living system - all that's
	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3127800 | 05/08/2024 | Copyright antibodies-online. All rights reserved.

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)