

Datasheet for ABIN3127835

WDR77 Protein (AA 1-342) (Strep Tag)



[Go to Product page](#)

Overview

| | |
|-------------------------------|------------------------------------------------|
| Quantity: | 1 mg |
| Target: | WDR77 |
| Protein Characteristics: | AA 1-342 |
| Origin: | Mouse |
| Source: | Cell-free protein synthesis (CFPS) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This WDR77 protein is labelled with Strep Tag. |
| Application: | ELISA, SDS-PAGE (SDS), Western Blotting (WB) |

Product Details

| | |
|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Brand: | AliCE® |
| Sequence: | <p>MRKDTPPPLV PPAAREWNLP PNAPACMERQ LEAARYRSDG SLLLGVSLSL GRCWVGSLWF FKDPSAAPNE GFCSAGVQTE AGVADLTWVG DKGILVASDS GAVELWELDE NETLIVSKFC KYEHDDIVST VTVLSSGTQA VSGSKDCCIK IWDLAQQVSL NSYRAHAGQV TCVAASPHKD SVFLSCSEDS RILLWDTRCP KPASQMACNA SGYLPTALAW HPQQSEVFVF GDENGVSLSLV DTKNASCTLS SAVHSQGVTR LVFSPHSVPL LTSLSLDCSL AVLDSSLSEV FRRAHRDFV RDATWSPLNH SLLTTVGWDH QVIHHVVPLE PLPNPGPDSV VE</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p> |
| Characteristics: | Key Benefits: |

Product Details

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

| | |
|---------------|--------------------------------------------------------------------------------------------------------------|
| Purification: | One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®). |
| Purity: | > 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC). |
| Grade: | custom-made |

Target Details

| | |
|---------|-------|
| Target: | WDR77 |
|---------|-------|

Target Details

| | |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alternative Name: | Wdr77 (WDR77 Products) |
| Background: | <p>Methylosome protein WDR77 (Methylosome protein 50) (MEP-50) (WD repeat-containing protein 77),FUNCTION: Non-catalytic component of the methylosome complex, composed of PRMT5, WDR77 and CLNS1A, which modifies specific arginines to dimethylarginines in several spliceosomal Sm proteins and histones (By similarity). This modification targets Sm proteins to the survival of motor neurons (SMN) complex for assembly into small nuclear ribonucleoprotein core particles (By similarity). Might play a role in transcription regulation (By similarity). The methylosome complex also methylates the Piwi proteins (PIWIL1, PIWIL2 and PIWIL4), methylation of Piwi proteins being required for the interaction with Tudor domain-containing proteins and subsequent localization to the meiotic nuage (PubMed:19584108). {ECO:0000250 UniProtKB:Q9BQA1, ECO:0000269 PubMed:19584108}., FUNCTION: Substrate-recognition component of the DCX(WDR77) complex, which mediates ubiquitination and degradation of Irgm1 in intestinal cells. {ECO:0000269 PubMed:35197566}.</p> |
| Molecular Weight: | 36.9 kDa |
| UniProt: | Q99J09 |
| Pathways: | Ribonucleoprotein Complex Subunit Organization |

Application Details

| | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. |
| Comment: | <p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p> |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Format: | Liquid |
| Buffer: | The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | 12 months |