Datasheet for ABIN3130282

FUT2 Protein (AA 1-347) (Strep Tag)

Go to Product page

Overview

Quantity:	1 mg
Target:	FUT2
Protein Characteristics:	AA 1-347
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This FUT2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)

Product Details

Sequence:	MASAQVPFSF PLAHFLIFVF VTSTIIHLQQ RIVKLQTLSE KELQAVQMSS PNAARTDMQQ SAKLQGIFTI NSIGRLGNQM GEYATLFALA RMNGRLAFIP ESMHNALAPI FRISLPVLHS DTARRIPWQN YHLNDWMEER YRHIPGQYVR FTGYPCSWTF YHHLRPEILK EFTLHDHVRE EAQAFLRGLR VNGSQPSTFV GVHVRRGDYV HVMPKVWKGV VADRGYLEKA LDRFRARYSS PVFVVTSNGM AWCRENINTS LGDVVFAGNG IEGSPAKDFA LLTQCNHTIM TIGTFGIWAA YLAGGDTIYL ANYTLPDSPF LKIFKPAAAF LPEWMGIPAD LSPLLKH **Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**
Characteristics:	Key Benefits: • Made in Germany - from design to production - by highly experienced protein experts. • Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure

correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.
The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®): 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details

Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

Target Details

Target:	FUT2
Alternative Name:	Fut2 (FUT2 Products)
Background:	Galactoside alpha-(1,2)-fucosyltransferase 2 (Alpha(1,2)FT 2) (Fucosyltransferase 2) (GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase 2) (GDP-fucose: beta-galactoside alpha1,2-fucosyltransferase) (MFUT-II) (Secretory blood group protein 2) (Type 1 galactoside alpha-(1,2)-fucosyltransferase FUT2) (EC 2.4.1.69) (Type 2 galactoside alpha-(1,2)-fucosyltransferase FUT2) (EC 2.4.1.344),FUNCTION: Catalyzes the transfer of L-fucose, from a guanosine diphosphate-beta-L-fucose, to the terminal galactose on both O- and N-linked glycans chains of cell surface glycoproteins and glycolipids and the resulting epitope regulates several processes such as cell-cell interaction including host-microbe interaction, cell surface expression and cell proliferation (PubMed:11018479, PubMed:11368156, PubMed:14967068, PubMed:11323419, PubMed:27161092, PubMed:19706747). Preferentially fucosylates gangliosides GA1 and GM1 in the antrum, cecum and colon and in the female reproductive organs (PubMed:11713270, PubMed:14967068). Fucosylated host glycoproteins or glycolipids mediate interaction with intestinal microbiota influencing its composition (PubMed:27161092, PubMed:19706747). Creates a soluble precursor oligosaccharide FuC-alpha ((1,2)Galbeta-) called the H antigen which is an essential substrate for the final step in the soluble ABO blood group antigen synthesis pathway (PubMed:11323419). {ECO:0000269|PubMed:11018479, ECO:0000269|PubMed:11323419, ECO:0000269|PubMed:11368156, ECO:0000269|PubMed:11713270, ECO:0000269|PubMed:14967068, ECO:0000269|PubMed:19706747, ECO:0000269|PubMed:27161092}.
Molecular Weight:	39.2 kDa
UniProt:	Q9JL27

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

Application Details

	Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)