antibodies .- online.com





CK1 epsilon Protein (AA 1-416) (His tag)



Image



Go to Product page

Overview

| Quantity: | 1 mg |
|-------------------------------|--|
| Target: | CK1 epsilon (CSNK1E) |
| Protein Characteristics: | AA 1-416 |
| Origin: | Mouse |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CK1 epsilon protein is labelled with His tag. |
| Application: | SDS-PAGE (SDS), ELISA, Western Blotting (WB), Crystallization (Crys) |

Product Details

Sequence:

MELRVGNKYR LGRKIGSGSF GDIYLGANIA SGEEVAIKLE CVKTKHPQLH IESKFYKMMQ
GGVGIPSIKW CGAEGDYNVM VMELLGPSLE DLFNFCSRKF SLKTVLLLAD QMISRIEYIH
SKNFIHRDVK PDNFLMGLGK KGNLVYIIDF GLAKKYRDAR THQHIPYREN KNLTGTARYA
SINTHLGIEQ SRRDDLESLG YVLMYFNLGS LPWQGLKAAT KRQKYERISE KKMSTPIEVL
CKGYPSEFST YLNFCRSLRF DDKPDYSYLR QLFRNLFHRQ GFSYDYVFDW NMLKFGAARN
PEDVDRERRE HEREERMGQL RGSATRALPP GPPTGATANR LRSAAEPVAS TPASRIQQTG
NTSPRAISRA DRERKVSMRL HRGAPANVSS SDLTGRQEVS RLAASQTSVP FDHLGK
Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Csnk1e Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

| Purity: | >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
|------------------|--|
| Sterility: | 0.22 μm filtered |
| Endotoxin Level: | Protein is endotoxin free. |
| Grade: | Crystallography grade |

Target Details

| Target: | CK1 epsilon (CSNK1E) |
|-------------------|--------------------------|
| Alternative Name: | Csnk1e (CSNK1E Products) |

Target Details

| rarget Details | |
|---------------------|--|
| Background: | Casein kinases are operationally defined by their preferential utilization of acidic proteins such |
| | as caseins as substrates. Can phosphorylate a large number of proteins. Participates in Wnt |
| | signaling. Phosphorylates DVL1. Central component of the circadian clock. In balance with PP1, |
| | determines the circadian period length, through the regulation of the speed and rhythmicity of |
| | PER1 and PER2 phospohorylation. Controls PER1 and PER2 nuclear transport and degradation. |
| | Inhibits cytokine-induced granuloytic differentiation. {ECO:0000269 PubMed:10848614, |
| | ECO:0000269 PubMed:14701732, ECO:0000269 PubMed:18400165, |
| | ECO:0000269 PubMed:19414593, ECO:0000269 PubMed:21930935}. |
| Molecular Weight: | 48.3 kDa Including tag. |
| UniProt: | Q9JMK2 |
| Pathways: | Hedgehog Signaling, M Phase |
| Application Details | |
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies |
| | as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee |
| | though. |
| Comment: | Protein has not been tested for activity yet. In cases in which it is highly likely that the |
| | recombinant protein with the default tag will be insoluble our protein lab may suggest a higher |
| | molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible |
| | options with you in detail to assure that you receive your protein of interest. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |
| | |



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process