

Datasheet for ABIN3130499

## NAA10 Protein (AA 1-235) (Strep Tag)



[Go to Product page](#)

### Overview

Quantity:	250 µg
Target:	NAA10 (ARD1A)
Protein Characteristics:	AA 1-235
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NAA10 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

### Product Details

Brand:	ALiCE®
Sequence:	<p>MNIRNARPED LMNMQHCNLL CLPENYQMKY YFYHGLSWPQ LSYIAEDENG KIVGYVLAKM  EEDPDDVPHG HITSLAVKRS HRRLGLAQKL MDQASRAMIE NFNAYVSLH VRKSNRAALH  LYSNTLNFQI SEVEPKYYAD GEDAYAMKRD LTQMADELRR HLELKEKGKH MVLAALENKA  ENKGNVLLSS GEACREEKGL AAEDSGGDSK DLSEVSETTE STDVKDSSEA SDSAS</p> <p><b>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>
Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"> <li>• Made in Germany - from design to production - by highly experienced protein experts.</li> <li>• Protein expressed with ALiCE® and purified in one-step affinity chromatography</li> <li>• These proteins are normally active (enzymatically functional) as our customers have</li> </ul>

## Product Details

---

reported (not tested by us and not guaranteed).

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

## Target Details

---

Target:	NAA10 (ARD1A)
Alternative Name:	Naa10 ( <a href="#">ARD1A Products</a> )
Background:	N-alpha-acetyltransferase 10 (EC 2.3.1.255) (N-terminal acetyltransferase complex ARD1

## Target Details

subunit homolog A) (NatA catalytic subunit Naa10),FUNCTION: Catalytic subunit of the N-terminal acetyltransferase A (NatA) complex which displays alpha (N-terminal) acetyltransferase activity (PubMed:12888564). Acetylates amino termini that are devoid of initiator methionine (By similarity). The alpha (N-terminal) acetyltransferase activity may be important for vascular, hematopoietic and neuronal growth and development (By similarity). Without NAA15, displays epsilon (internal) acetyltransferase activity towards HIF1A, thereby promoting its degradation (PubMed:12464182). Represses MYLK kinase activity by acetylation, and thus represses tumor cell migration (By similarity). Acetylates, and stabilizes TSC2, thereby repressing mTOR activity and suppressing cancer development (By similarity). Acetylates HSPA1A and HSPA1B at 'Lys-77' which enhances its chaperone activity and leads to preferential binding to co-chaperone HOPX (By similarity). Acetylates HIST1H4A (By similarity). Acts as a negative regulator of sister chromatid cohesion during mitosis (By similarity). {ECO:0000250|UniProtKB:P41227, ECO:0000269|PubMed:12464182, ECO:0000269|PubMed:12888564}.

Molecular Weight: 26.5 kDa

UniProt: [Q9QY36](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months