antibodies .- online.com





ITGB1BP1 Protein (AA 1-200) (Strep Tag)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	1 mg
Target:	ITGB1BP1
Protein Characteristics:	AA 1-200
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ITGB1BP1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Seqı	Jence:
------	--------

MFRKGKKRHS SSSSQSSEIS TKSKSVDSSL GGLSRSSTVA SLDTDSTKSS GQSNSNLDTC
AEFRIKYVGA IEKLAVSEGK SLEGPLDLIN YIDVAQQDGK LPFVPLEEEF ILGVSKYGIK
VSTTDQHGVL HRHALYLIIR MVCYDDGLGA GKSLLALKTT DASNEEYSLW VYQCNSLEQA
QAICKVLSTA FDSVLTSDKS

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- · These proteins are normally active (enzymatically functional) as our customers have

reported (not tested by us and not guaranteed).

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Target Details

Target:	ITGB1BP1	
Alternative Name:	Itgb1bp1 (ITGB1BP1 Products)	
Background:	Integrin beta-1-binding protein 1 (Bodenin),FUNCTION: Key regulator of the integrin-mediated	
	cell-matrix interaction signaling by binding to the ITGB1 cytoplasmic tail and preventing the	
	activation of integrin alpha-5/beta-1 (heterodimer of ITGA5 and ITGB1) by talin or FERMT1.	
	Plays a role in cell proliferation, differentiation, spreading, adhesion and migration in the contex	
	of mineralization and bone development and angiogenesis. Stimulates cellular proliferation in a	
	fibronectin-dependent manner. Involved in the regulation of beta-1 integrin-containing focal	
	adhesion (FA) site dynamics by controlling its assembly rate during cell adhesion, inhibits beta-	
	1 integrin clustering within FA by directly competing with talin TLN1, and hence stimulates	
	osteoblast spreading and migration in a fibronectin- and/or collagen-dependent manner. Acts	
	as a guanine nucleotide dissociation inhibitor (GDI) by regulating Rho family GTPases during	
	integrin-mediated cell matrix adhesion, reduces the level of active GTP-bound form of both	
	CDC42 and RAC1 GTPases upon cell adhesion to fibronectin. Stimulates the release of active	
	CDC42 from the membranes to maintain it in an inactive cytoplasmic pool. Participates in the	
	translocation of the Rho-associated protein kinase ROCK1 to membrane ruffles at cell leading	
	edges of the cell membrane, leading to an increase of myoblast cell migration on laminin. Plays	
	a role in bone mineralization at a late stage of osteoblast differentiation, modulates the	
	dynamic formation of focal adhesions into fibrillar adhesions, which are adhesive structures	
	responsible for fibronectin deposition and fibrillogenesis. Plays a role in blood vessel	
	development, acts as a negative regulator of angiogenesis by attenuating endothelial cell	
	proliferation and migration, lumen formation and sprouting angiogenesis by promoting AKT	
	phosphorylation and inhibiting ERK1/2 phosphorylation through activation of the Notch	
	signaling pathway. Promotes transcriptional activity of the MYC promoter.	
	{ECO:0000269 PubMed:16741948, ECO:0000269 PubMed:17567669,	
	ECO:0000269 PubMed:17654484, ECO:0000269 PubMed:18227284,	
	ECO:0000269 PubMed:21768292}.	
Molecular Weight:	21.6 kDa	
UniProt:	035671	
Pathways:	Tube Formation	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	

Application Details

- 1 1		
	as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from	
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce	
	even the most difficult-to-express proteins, including those that require post-translational	
	modifications.	
	During lysate production, the cell wall and other cellular components that are not required for	
	protein production are removed, leaving only the protein production machinery and the	
	mitochondria to drive the reaction. During our lysate completion steps, the additional	
	components needed for protein production (amino acids, cofactors, etc.) are added to produce	
	something that functions like a cell, but without the constraints of a living system - all that's	
	needed is the DNA that codes for the desired protein!	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request,	
	please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	