# antibodies -online.com





# COPS5 Protein (AA 2-334) (His tag)



#### Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	1 mg
Target:	COPS5
Protein Characteristics:	AA 2-334
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This COPS5 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

### **Product Details**

## Sequence:

AASGSGMAQK TWELANNMQE AQSIDEIYKY DKKQQQEILA AKPWTKDHHY FKYCKISALA LLKMVMHARS GGNLEVMGLM LGKVDGETMI IMDSFALPVE GTETRVNAQA AAYEYMAAYI ENAKQVGRLE NAIGWYHSHP GYGCWLSGID VSTQMLNQQF QEPFVAVVID PTRTISAGKV NLGAFRTYPK GYKPPDEGPS EYQTIPLNKI EDFGVHCKQY YALEVSYFKS SLDRKLLELL WNKYWVNTLS SSSLLTNADY TTGQVFDLSE KLEQSEAQLG RGSFMLGLET HDRKSEDKLA KATRDSCKTT IEAIHGLMSO VIKDKLFNOI NVA

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Cops5 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

## **Target Details**

Target:	COPS5
Alternative Name:	Cops5 (COPS5 Products)
Background:	Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in

various cellular and developmental processes. The CSN complex is an essential regulator of the
ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of the
SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type
complexes such as SCF, CSA or DDB2. Promotes the proteasomal degradation of BRSK2. The
complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, lkappaBalpha/NFKBIA,
ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent
phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system,
respectively. In the complex, it probably acts as the catalytic center that mediates the cleavage
of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the
other subunits of the CSN complex. Interacts directly with a large number of proteins that are
regulated by the CSN complex, confirming a key role in the complex.

Molecular Weight:	38.4 kDa Including tag.
UniProt:	O35864
Pathways:	Cell Division Cycle

## **Application Details**

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
	though.	
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the	
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher	
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible	
	options with you in detail to assure that you receive your protein of interest.	

For Research Use only

# Handling

Restrictions:

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

$\vdash$	land	linc
	iaria	11116

Expiry Date:

Unlimited (if stored properly)