Datasheet for ABIN3132030

SRPK1 Protein (AA 1-648) (Strep Tag)

Go to Product page

Overview

Quantity:	250 µg
Target:	SRPK1
Protein Characteristics:	AA 1-648
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SRPK1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	AliCE®
Sequence:	MERKVLALQA RKKRTKAKKD KAQRKPETQH RGSAPHSESD IPEQEEEILG SDDDEQEDPN DYCKGGYHLV KIGDLFNGRY HVIRKLGWGH FSTVWLSWDI QGKKFVAMKV VKSAEHYTET ALDEIRLLKS VRNSDPNDPN GEMVVQLLDD FKISGVNGTH ICMVFEVLGH HLLKWIIKSN YQGLPLPCVK KIIQQVLQGL DYLHTKCRII HTDIKPENIL LSVNEQYIRR LAAEATEWQR SGAPPPSGSA VSTAPQPKPA DKMSKNKKKK LKKKQKRQAE LLEKRMQEIE EMEKESGPGQ KRPNKQEESE SPVDRPLTEN PPNKMTQEKL EESNSIGQDQ TLTERGGEGG APEINCNGVI GVVNYPENSN NETLRHKEDL HNANDCDVHT LKQEPSFLNS SNGDSSPSQD TDSCTPTASE TMVCQSSAEQ SLTRQDITQL EESIRADTPS GDEQEPNGAL DSKGKFSAGN FLINPLEPKN AEKLQVKIAD LGNACWVHKH FTEDIQTRQY RSLEVLIGSG YNTPADIWST ACMAFELATG DYLFEPHSGE DYTRDEDHIA LIIELLGKVP RKLIVAGKYS KEFFTKKGDL KHITKLKPWG LLEVLVEKYE WPQEEAAGFT DFLLPMLELM PEKRATAAEC LRHPWLNS

Product Details

	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.
Characteristics:	Key Benefits: • Made in Germany - from design to production - by highly experienced protein experts. • Protein expressed with ALiCE® and purified in one-step affinity chromatography • These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed). • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. Expression System: • ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. • During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein! Concentration: • The concentration of our recombinant proteins is measured using the absorbance at 280nm. • The protein's absorbance will be measured against its specific reference buffer. • We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.
Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Product Details

Grade:	custom-made

Target Details

Target:	SRPK1
Alternative Name:	Srpk1 (SRPK1 Products)
Background:	SRSF protein kinase 1 (EC 2.7.11.1) (SFRS protein kinase 1) (Serine/arginine-rich protein-specific kinase 1) (SR-protein-specific kinase 1),FUNCTION: Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing. Plays a central role in the regulatory network for splicing, controlling the intranuclear distribution of splicing factors in interphase cells and the reorganization of nuclear speckles during mitosis. Can influence additional steps of mRNA maturation, as well as other cellular activities, such as chromatin reorganization in somatic and sperm cells and cell cycle progression. Phosphorylates SFRS2, ZRSR2, LBR and PRM1. Phosphorylates SRSF1 using a directional (C-terminal to N-terminal) and a dual-track mechanism incorporating both processive phosphorylation (in which the kinase stays attached to the substrate after each round of phosphorylation) and distributive phosphorylation steps (in which the kinase and substrate dissociate after each phosphorylation event). The RS domain of SRSF1 binds first to a docking groove in the large lobe of the kinase domain of SRPK1. This induces certain structural changes in SRPK1 and/or RRM2 domain of SRSF1, allowing RRM2 to bind the kinase and initiate phosphorylation. The cycles continue for several phosphorylation steps in a processive manner (steps 1-8) until the last few phosphorylation steps (approximately steps 9-12). During that time, a mechanical stress induces the unfolding of the beta-4 motif in RRM2, which then docks at the docking groove of SRPK1. This also signals RRM2 to begin to dissociate, which facilitates SRSF1 dissociation after phosphorylation is completed. Can mediate hepatitis B virus (HBV) core protein phosphorylation. It plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgRNA) without affecting the formation of the viral core particles. Can induce splicing of exon 10 in MAPT/TAU (By similarity). {ECO:0000250, ECO:0000269|PubMed:10390541, ECO:0000269|PubMed:9446799}.
Molecular Weight:	73.1 kDa
UniProt:	O70551
Pathways:	Toll-Like Receptors Cascades

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months