

Datasheet for ABIN3132220 STAM2 Protein (AA 1-523) (Strep Tag)



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	1 mg
Target:	STAM2
Protein Characteristics:	AA 1-523
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This STAM2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence:

MPLFTANPFE QDVEKATNEY NTTEDWSLIM DICDRVGSTP SGAKDCLKAI MKRVNHKVPH
VALQALTLLG ACVANCGKIF HLEVCSRDFA TEVRSVIKNK AHPKVCEKLK SLMVEWSEEF
QKDPQFSLIS ATIKSMKEEG VTFPSAGSQT VAAAAKNGTS LNKNKEDEDI AKAIELSLQE
QKQQYTETKA LYPPAESQLN NKAARRVRAL YDFEAVEDNE LTFKHGELIT VLDDSDANWW
QGENHRGTGL FPSNFVTTDL STEVETATVD KLNVIDDDVE EIKKSEPEPV YIDEGKMDRA
LQILQSIDPK ESKPDSQDLL DLEDVCQQMG PMIDEKLEEI DRKHSELSEL NVKVLEALDL
YNKLVNEAPV YSVYSKLHPA HYPPAAAGVP VQTYPVQSHG GNYLGHGIHQ VSVAQNYNLG
PDPMGSLRSL PPNMNSVTAH TVQPPYLSTG QDTVSNPSYM NQSSRLQAAA GTAAYTQPVG
MSTDVSSFQN TASGLPQLAG FPVAVPAPVA AQPQASYHQQ PLL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

Product Details

	Western blot.		
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.		
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)		
Target Details			
Target:	STAM2		
Alternative Name:	Stam2 (STAM2 Products)		
Background:	Signal transducing adapter molecule 2 (STAM-2) (Hrs-binding protein),FUNCTION: Involved in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSL stimulation, it plays a role in signaling leading to DNA synthesis and MYC induction. May also play a role in T-cell development. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with HGS (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as a sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes (By similarity). {ECO:0000250, ECO:0000269 PubMed:12972556}.		
Molecular Weight:	57.5 kDa		
UniProt:	088811		
Pathways:	EGFR Signaling Pathway, EGFR Downregulation		
Application Details			
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.		
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce		

Application Details

	needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)